

which it is more nearly allied, in the perfectly straight outer edge of the intermediate tibiae. The cicatrice of the scape is visible only in a certain light, and recognizable chiefly by its limiting margin; it is very broad, as in *Xenolea*.

NANOHAMMUS RUFESCENS, n. sp. (Plate II. fig. 4.)

Fulvo-castaneus, subnudus, thoracis linea dorsali, scutello maculisque elytrorum (ante et post medium subfasciatim congregatis) ochraceo-tomentosis; antennarum articulis 3-4 basi tibiisque pallidioribus; capite omnino discrete punctulato; thorace creberrime subconfluenter, elytris passim discrete hic illuc seriatim, punctatis. Long. 7 millim.

Chiuzenji.

SCOTINAUGES DIPHYSIS, *Pascoe*, *Ann. & Mag. N. H.* 1871, p. 277, t. xiii. fig. 4.

Island of Tsu-shima (*Bowring*).

This species was omitted from my first enumeration of the Longicornia of Japan.

MESOSA GRACILLIOR, n. sp.

M. nebulosæ (Oliv.) primo intuitu similis, sed angustior, gracilis, thorace præcipue angustiori, minori lateribusque rotundatis. Anguste oblonga, fulvescenti-fusco adpresso-pubescentes, vertice et thorace nigro pluri-maculatis (nec vittatis); elytris guttis sat confertis nigris fasciaque mediana dentata cinerea nigro partim marginata; antennis nigro-fuscis, articulis basi griseis; elytris elongato-oblongis, crebrius quam in *M. nebulosa* punctulatis. Long. 10-11 millim. ♂ ♀.

Oyayama.

Narrower and more elongated than *M. nebulosa*, the thorax especially relatively much smaller and narrower and rounded on the sides.

MESOSA JAPONICA, *Bates*.

S. Japan; also Yezo.

✓ **MESOSA HIRSUTA**, n. sp.

Elongato-oblonga, supra dense eructe pubescens, fusco-tomentosa, griseo plagiatum varia; thoracis maculis novem (interdum indistinctis), elytrorum maculis circa 20, quarum 4 suturalibus communibus, atro-fuscis; antennis fusco-nigris, articulis 3-11

basi griseis, scapo griseo et fusco-piperito, art. undecimo prædenti ♂ paullo, ♀ dimidio, breviori; thorace vix inæquali, lateribus leviter rotundatis, sparsim punctulato; elytris subtilius quam in *M. nebulosa* punctulatis; pedibus griseis nigro-annulatis sicut in *M. nebulosa*. Long. 12-16 millim. ♂ ♀.

Kobé.

MESOSA SENILIS, n. sp.

M. nebulosæ quoad formam similis, sed paullo magis elongata, canescenti-griseo tomentosa, thoracis vittis duabus angustis (sæpe obsoletis), elytrorum vitta curvata subhumerali, macula longe post medium marginali maculisque suturalibus discoidalibusque (sæpe deficientibus) nigra. Antennis griseis, articulis 3-10 apice breviter infuscatis, ♂ corpore dimidio longioribus, articulo undecimo ♂ elongato precedenti sublongiori, ♀ præcedenti breviori; thorace sicut in *M. nebulosa* inæquali; scutello toto griseo; elytris sparsim punctatis, punctis nigra. Long. 11-12 millim. ♂ ♀.

Junsai; Oyama; Sapporo.

Varies much in the number and distinctness of the black spots and marks; the hoary laid pubescence is sometimes thin and coarse, and then seems to cover the black marks. In some examples the prevailing colour is light brownish-grey instead of light grey.

MESOSA PÆCILA, n. sp.

Elongato-oblonga, eructe nigro-pubescentes, griseo-vel ochraceo-griseo tomentosa tota nigro-piperita, fascia undulata elytrorum longe post medium (sutura haud attingenti) maculisque 2-3 marginalibus prope humerum, nigra. Capite inter antennas profundius quam in *M. nebulosa* concavo; thorace relative parvo, juxta basin angustato, dorso haud tuberculato sed antice transversim depresso; scutello griseo, lateribus nigra; elytris punctulatis; antennis griseis, articulis omnibus apice nigra, ♀ (?) art. undecimo precedenti dimidio breviori; pedibus griseis nigro-maculatis. Long. 13-17 millim.

Nikko; Junsai.

MESOSA CRIBRATA, n. sp.

Parva, oblonga, eructe pilosa, griseo-fusca tomentosa, elytrorum fasciis duabus latis (altera recta basali, altera post mediana undulata) atrofuscis. Capite inter antennas parum concavo; thorace brevi cylindrico, supra æquali, sparsim grosse punctato;