



The geographic range of *Tragosoma depsarium* (Linnaeus, 1767) (Coleoptera, Cerambycidae) in the Palaearctic

Nikolay S. Anisimov^{1*}, Vitaly G. Bezborodov²

1 Federal State Budget Scientific Institution Federal Scientific Center All-Russian Scientific Research Institute of Soybean, Blagoveshchensk, Russia • havamall@mail.ru

2 Amur Branch of Botanical Garden-Institute, Far Eastern Branch, Russian Academy of Sciences, Blagoveshchensk, Russia • cichrus@yandex.ru

* Corresponding author

Abstract

The geographic range of the rare coleopteran *Tragosoma depsarium* (Linnaeus, 1767) is summarized, and the Eurasian distribution is mapped for the first time. Particular attention is paid to the eastern Palaearctic part of the species' range, where dubious data prevail. Distributional data from the Russian Far East have been published, but not confirmed, are considered, and new records of this species in Eastern Siberia are reported.

Keywords

Distribution, Europe, rare species, Russian Far East, Siberia

Academic editor: Cheng-Bin Wang | Received 4 February 2021 | Accepted 23 May 2021 | Published 31 May 2021

Citation: Anisimov NS, Bezborodov VG (2021) The geographic range of *Tragosoma depsarium* (Linnaeus, 1767) (Coleoptera, Cerambycidae) in the Palaearctic. Check List 17 (3): 841–851. <https://doi.org/10.15560/17.3.841>

Introduction

Tragosoma depsarium (Linnaeus, 1767) is the sole species of the genus *Tragosoma* Audinet-Serville, 1832 and the tribe Meroscelisini J. Thomson, 1861 in the Palaearctic (Drumont and Komiya 2010). Previously, this species was considered Holarctic; however, modern data show that it does not occur outside the Palaearctic (Laplante 2017).

Throughout its entire range, *T. depsarium* is rarely and sporadically found. This species is ecologically vulnerable. Rapid reduction of forests with optimal habitats is the main limiting factor for this species' distribution. *Tragosoma depsarium* prefers coniferous forests where the proportion of mature trees is more than 25%, lighting conditions are good and stand density is low, and there is considerable dry, windfallen pines or spruces having a large diameter and no bark. Soil under these trees should

have an optimal moisture content, which is important for symbiotic fungi (Wikars 2003, 2004). The current forest management regulations, including extinguishing forest fires, sanitary felling of old trees and clearing of wind-fallen trees, as well as replacing mature forests cut for industrial purposes with man-made, high-density stands, do not ensure good environmental conditions for this rare species. Even within protected areas, where stand density becomes too high without felling and fires, populations of this species can be threatened. The gaps between the isolated populations also impair the chances of this species' long-term survival.

Tragosoma depsarium is protected in Europe (Nieto and Alexander 2010) and many regions of Russia (Jakovlev 2007; Tikhomirov 2007; Nikitsky 2008; Egorov 2010; Dedyukhin 2012; Sitnikov 2013; Potanin 2014; Yuferev

2014; Vlasov 2015; Gus'kova 2016; Gorbunov 2018; Shapovalov 2018; Dolgin and Kolesnikova 2019). However, more accurate data on this species' distribution are needed to ensure efficient protective measures. It is not always possible to be sure that *T. depsarium* still occurs in sites where it was found over 100 years ago, and its presence has not been confirmed later. The fact that vast forests are now growing in places where old forests used to grow does not guarantee that *T. depsarium* population can be where they once were. Because of the strict ecological requirements of *T. depsarium* and the possible changes in local conditions, this species could have disappeared in these parts of the range. Meanwhile, records have recently been made for the first time in some territories. The data on the distribution of *T. depsarium* in the Asian part of Russia are insufficient. In this study, we attempt to summarize the scattered data on the distribution of *T. depsarium* from all available sources. We also present new distributional data from the eastern part of the species' range.

Methods

We used an extensive list of available literary sources, web resources, and personal reports from observers and collectors to summarize the geographic range of *Tragosoma depsarium*. The entomological collections of the Zoological Institute of the Russian Academy of Sciences (Saint Petersburg), Federal Scientific Center for Terrestrial Biodiversity of East Asia FEB RAS (Vladivostok)

and the Institute of Systematics and Ecology of Animals SB RAS (Novosibirsk) were studied. Furthermore, we publish for the first time data on the easternmost record of *T. depsarium*, which we confirmed by the study of the actual material.

The image of the specimens from Zabaykalsky Krai was taken using a Canon EOS 5d Mark II camera with a Volna-9 50 mm f/2.8 macro lens. The maps were prepared using the CorelDRAW Graphics Suite X7 software.

The following symbols were used in the text and on the map: in the text – Tomsk Oblast (TOM): Pervomaysky District (3); on the map – TOM3. No figure denotes that no other records were made within this territory: ZAB – Zabaykalsky Krai. The material provided by the Institute of Systematics and Ecology of Animals is referred to as ISEA in the text; Zoological Institute of the Russian Academy of Sciences – ZIN. Data on the age of the records are graphically shown on the maps.

Results

Tragosoma depsarium (Linnaeus, 1767)

New record. RUSSIA – Zabaykalsky Krai • Shilkinsky District, 7 km S of Kyeken village, Ukshakhan creek; 52°00'00"N, 115°25'08"E; 602 m elev.; 30.VI.2017; V.A. Golovizin leg.; small-leaved forest; 1♀, 1♂, V.G. Bezborrowodov's personal collection (Fig. 1).




Figure 1. *Tragosoma depsarium* (dorsal view) from Zabaykalsky Krai, Russia. **A.** Male. **B.** Female.

Additional material examined. UKRAINE – **Rivne Oblast** • Rivne, 1898; 1 specimen, ZIN. RUSSIA – **Arkhangelsk Oblast** • Velsk; D.V. Pomerantsev leg.; 1 specimen, ZIN. **Chelyabinsk Oblast** • Zlatoust, 4.VIII.1898; Kalachev leg.; 1 specimen, ZIN. **Tomsk Oblast** • Pervomaysky District, Frantsevo village, 6.VII.1959; Grigoriev leg.; 1 specimen, ISEA • mouth of the Chichkayul River, 18.VI-8.VII.1959; 2 specimens, ISEA. **Novosibirsk Oblast** • Novosibirsk (“Novonikolayevsk”), 22.VI.1910; Kopylov leg.; 1 specimen, ZIN • Bolshaya Elovka River, 1.VIII.1912; A. Mylenikoff leg.; 3 specimens, ZIN. **Kemerovo Oblast** • Novokuznetsky District, Kalinovskiy village, VII.1979; 1 specimen, ISEA • Tashtagolsky District, Sheregesh settlement, 9.VII.1965; A. Kononenko leg.; 2 specimens, ISEA. **Altai Krai** • Troitsky District, vicinity of Borovlyanka village, 3.VII.1953; 1 specimen, ISEA. **Altai Republic** • Ulagansky District, S of Lake Teletskoye, 10.VII.1969; 3 specimens, ISEA • Choysky District, 20 km S of Paspaul village, 20.VII.1999; V.K. Zinchenko leg.; 1 specimen, ISEA • Shebalinsky District, Cherginsky Ridge, the Kukuya River, early July 1981; V.V. Dubatolov leg.; 1 specimen, ISEA • Chemalsky District, Uznezya village, 29.VI.1909; Gorchakovskiy leg.; 1 specimen, ZIN. **Republic of Khakassia** • Askizsky District, Birikchul village, 25.VII.1969; 1 specimen, ISEA. **Krasnoyarsk Krai** • vicinity of Krasnoyarsk, VIII.1901; 1 specimen, ZIN.

Identification. *Tragosoma depsarium*, the only of the genus *Tragosoma* and tribe Meroscelisini in the Palaearctic fauna, morphologically differs very much from other species of this subfamily found in Eurasia, which makes it is easily identifiable. The imago has elongated elytra and a small pronotum and is brownish-rusty in color. The head is small, with a groove between eyes. The antennae have the apices reaching the middle of the elytra in males, or not reaching this level in females. The pronotum bears dense, rusty hairs. The body is brownish-rusty and ventrally has long, golden hairs.

Discussion

The range of *Tragosoma depsarium* can be conditionally subdivided into two parts: the European one, which includes the European countries and Russia stretching as far as the Ural Mountains, and the Asian one, including only Siberia, where the easternmost limits of the species' range are insufficiently known. We summarize the distributional data of *T. depsarium* and the conservation status of the species in various territories, as well as present maps showing the locations and the age of the records.

The European part of the range, European countries (Fig. 2). **Norway (NR):** Aust-Agder (1), Telemark (2), Vestfold (3), Buskerud (4), Hedmark (5) (Blindheim 2020; Ødegaard and Vang 2020). Conservation status: Vulnerable (Ødegaard et al. 2015). **Sweden (SV):** Lappland (1), Norrbotten (2), Västerbotten (3), Ångermanland (4), Jämtland (5), Medelpad (6), Härjedalen (7),

Hälsingland (8), Gästrikland (9), Dalarna (10), Värmland (11), Västmanland (12), Närke (13), Uppland (14), Södermanland (15), Dalsland (16), Östergötland (17), Småland (18), and Gotska Sandön National Park (19) (Wikars 2003; GBIF 2020); previously in Västergötland (Thomson 1866). Conservation status: Vulnerable in 2010 (Ljungberg et al. 2010) but changed to Near Threatened in 2015 (Ljungberg 2015). **Finland (FI):** North Karelia (1) since 1975; earlier in Pohjois-Savo (2), Pirkanmaa (3), Keski-Suomi (4), Kanta-Häme (5), Varsinais-Suomi (6) (Saalas 1923; Karjalainen et al. 2010); Etelä-Pohjanmaa (South Ostrobothnia) (Saalas 1923). Conservation status: Endangered (Malmberg et al. 2019). **Spain (SP):** Huesca (1), Lleida (2) (González et al. 2007). **Andorra (AN):** Botella Pass in 1975 (Vives 1984). **France (FR):** Upper Pyrenees (1), Upper Garonne (2), Eastern Pyrenees (3), Upper Savoy (4), Savoy (5), Maritime Alps (6) (Brin et al. 2006; Brustel et al. 2001; Blanc 2011; Turpaud 2018; Plaige 2019; Duhem and Horellou 2020; Lapenu 2020; Touroult 2020). **Belgium (BE):** old records near Brussels (1) and Chaudfontaine (2) (Verschelde and Brosens 2019a, 2019b). **Germany (GE):** mountain forests of Upper Bavaria (1) (Müller-Kroehling et al. 2009; European Nucleotide Archive 2019); eastern Thuringia (2) in 2005, where it had probably been introduced (Flossmann 2006); included in the lists of threatened species of Germany and Bavaria: Endangered (Binot et al. 1998; Schmidl and Bussler 2003). **Switzerland (SZ):** Vaud (1), Fribourg (2), Valais (3), Ticino (4) (Chittaro et al. 2020). Conservation status: Endangered (Monnerat et al. 2016). **Austria (AU):** Vorarlberg in 2016 (1) (Eckelt and Degasperi 2018); North Tyrol (2), Salzburg (3), Styria (4), Upper Austria (5), Lower Austria (6), Carinthia (7) (Demelt 1957; Lovcik 1970; Mitter 1988; Schillhammer 1995; Adlbauer 2001, 2010; Hellrigl 2010; Kust 2016; Aurenhammer et al. 2019). Conservation status: possibly Endangered (Jäch 1994). **Liechtenstein (LS):** repeated records in this country (Adlbauer 1992; Brandstetter and Kapp 1998; Drumont and Komiya 2010; Mannerkoski et al. 2010). **Italy (IT):** 19th century records at Great St. Bernard Pass, Piedmont (1) (Ghiliani 1887), Val Cadino, province of Trento (2) (Bertolini 1895); South Tyrol (3) (Bertolini 1895; Hellrigl 1967); Val Genova, province of Trento (Moscardini 1956), have not been subsequently confirmed (Martinelli 1996). Conservation status: Critically Endangered (Rapuzzi et al. 2014). **Poland (PL):** last found in Białowieża Forest (Belovezhskaya Pushcha) (1) in 1921; before that, found in the Warmian-Masurian (2), Kuyavian-Pomeranian (3) and Lower Silesian (4) provinces (Gutowski 2004). Conservation status: Critically Endangered (Pawłowski et al. 2002). **Czech Republic (CZ):** South Bohemia (1), Moravian-Silesian Beskids (2). Until 1949, in the Central Bohemian Region (3) (Sláma 1998; Krása 2015). Conservation status: Critically Endangered (Kabátek and Skořepa 2017). **Slovakia (SK):** Žilina (1), Prešov (2) and Banská Bystrica (3) regions (Sláma 1998). Conservation status: Vulnerable (Holecová and Franc 2001). **Hungary (HU):** a single



Figure 2. Distribution records of *Tragosoma deparium* (Linnaeus, 1767) in Europe.

specimen in the Zemplén Mountains in north-eastern Hungary in 1986 (Medvegy 2001). **Slovenia (SL)**: one specimen in the Julian Alps in northern Gorizia Region (1) in 1952; additional 19th century records in Carniola (2) (Breljih et al. 2006). **Croatia (CR)**: Velika Kapela (1) and Velebit mountain ranges in the western Croatia; Virovitica-Podravina County (2) (Mikšić and Georgijević 1971); Nikola Rahme (pers. comm.) reported that he found larvae and adults in the Northern Velebit National

Park (3) in 2007. **Serbia (SB)**: Kopaonik Mountains in 1911 (1) (Ilić and Ćurčić 2015; Mesaroš 2020); according to Filip Vukajlović and Dragan Pavićević (pers. comms.), in Kosovo (UTM DN31) on 23 August 1918, in the Prokletije Mountains (2). In the summer of 1918, this region of Kosovo constituted a part of the Kingdom of Montenegro, so in the literature (Mikšić 1963; Mikšić and Georgijević 1971) reportedly made in Montenegro. **Bosnia and Herzegovina (BH)**: Federation of Bosnia and

Herzegovina: Glamoč (1), Livno (2), Sarajevo (3), Mount Igman (4); Republika Srpska: Bijelo Brdo (5) (Apfelbeck 1894; Mikšić 1963). **Albania (AL)**: Shkodër (1) and Dibër (2) (Foit 2007; Rapuzzi and Sama 2012). **Greece (GR)**: Katara Pass in the Pindus Mountains (1), Mount Olympus (2) (Berger et al. 2010). **Bulgaria (BU)**: Pazardzhik (1), Blagoevgrad (2) and Smolyan (3) provinces (Migliaccio et al. 2007). Conservation status: Vulnerable (Guéorguiev 2015). **Romania (RO)**: Transylvania, Harghita Mountains (Patkó 2008). **Estonia (EN)**: extirpated (Süda 2008). More than a century ago, it was found near Tartu (1) and north of Lake Peipsi, now within modern Ida-Viru County (2) (Voolma and Süda 1999). **Latvia (LA)**: extirpated (Spuris 1998), but in 1999–2002, fragments of adults in Valka (1), Alūksne (2), and Ventspils (3) municipalities (Telnov and Kalniņš 2003); Slitere National Park and its surroundings (4) in 2008–2019 (Vilks and Pilāts 2008; Dabasdati.lv 2020); Gulbene Municipality (5) (Telnov et al. 2011). **Lithuania (LT)**: vicinity of Vilnius (1) in the 1920s (Zawadzki 1937); Chepkeliai Nature Reserve (2) in 1978 (Pileckis and Monsevičius 1997). **Belarus (BY)**: Vitebsk (1), Mogilev (2), Brest (3) regions; Belovezhskaya Pushcha National Park (4) (Arnold 1902; Radkevich 1936; Zagaykevich 1986; Tsinkevich 2015). Conservation status: Endangered. **Ukraine (UA)**: Eastern Beskids (1) (Nowicki 1864; Zamoroka 2018); Olevsk (2) (Plavilstshikov 1936); very old records in the vicinity of Kyiv (3) (Cherkunov 1889) and Rivne (4) (additional material, ZIN).

The European part of the range, Russia (Fig. 3). Republic of Karelia (KR): Suoyarvsky (1), Kondopozhsky (2),

Olonetsky (3), Pudozhsky (4) and Pitkyarantsky (5) districts, Kostomukhsky Urban Okrug (6) (Saalas 1923; Siitonen et al. 1996; Jakovlev et al. 2003; Humala and Polevoi 2009). Conservation status: Endangered (Jakovlev 2007). **Arkhangelsk Oblast (ARK)**: vicinity of Velsk before 1905 (additional material, ZIN). **Leningrad Oblast (LEN)**: Vyborgsky (1), Priozersky (2), Lodeynopolsky (3) Luzhsky (4) districts (Saalas 1923; Krivokhatsky 2002; Shapovalov 2018); old records from the former Terijoki, now within St. Petersburg (5) (Saalas 1923). Conservation status: category 4 – undefined status (Shapovalov 2018). Previous conservation status: Vulnerable (Krivokhatsky 2002). **Pskov Oblast (PSK)**: vicinity of Pskov in 1900 (Chistovsky 1929). **Bryansk Oblast (BRY)**: Bryansk forest at the beginning of the 20th century (Stark 1926). **Kaluga Oblast (KLU)**: probably extirpated (Alekseev and Perov 2017); Lyudinovo forest at the beginning of the 20th century (Chernyshov 1930). **Moscow Oblast (MOS)**: Solnechnogorsky, Istrinsky and Odintsovsky districts (1) over 50 years ago; Orekhovo-Zuyevesky District (2) in 2006 (Danilevsky 2005; Nikitsky 2008; Nikitsky 2009). Conservation status: rare species declining in number. **Yaroslavl Oblast (YAR)**: Rybinsky, Uglichsky (1), Pereslavsky (2) districts (Vlasov 2019); vicinity of Yaroslavl (3) in the 19th century (Vlasov 2015). Conservation status: Endangered (Vlasov 2015). **Ivanovo Oblast (IVA)**: Kineshemy (1), Pestyakovsky, and Yuzhsky (2) districts (Svetlov 1996; Danilevsky 2014). Conservation status: rare species (Tikhomirov 2007). **Nizhny Novgorod Oblast (NIZ)**: Volodarsky District (1), Semyonovsky Urban Okrug (2) (Mokrousov 2008). Conservation status: species with




Figure 3. Distribution records of *Tragosoma depsarium* (Linnaeus, 1767) in Russia.

low abundance (population density) being the biological norm (Potanin 2014). **Kirov Oblast (KIR)**: Shabalinsky, Svechinsky (1), Kotelnichsky (2), and Nolinsky (3) districts. Conservation status: rare species declining in number (Yuferev 2014). **Mari El Republic (ME)**: one specimen in the Zvenigovsky District in 1989 (Ivshin 1990). **Chuvash Republic (CU)**: Zavolzhye (near Cheboksary) (1); Alatyrsky and Shemurshinsky (2) districts (Egorov 2009, 2018). Conservation status: Rare species not threatened with extinction (Egorov 2010). **Komi Republic (KO)**: vicinity of Syktyvkar, Syktyvdinsky (1); Sysolsky (2), Kortkerossky districts (3); upper reaches of the Pechora River (4) (Sedykh 1974; Medvedev 1997; Yurkina 2011; Dolgin and Kolesnikova 2019). Conservation status: rare species with naturally low abundance (Dolgin and Kolesnikova 2019). **Udmurt Republic (UD)**: Yakshur-Bodyinsky (1), Zavyalovsky (2), Sarapulsky (3), Kambarsky (4) districts (Dedyukhin 2005, 2006, 2007; Adakhovsky and Dedyukhin 2011; Baranova et al. 2011). Conservation status: rare species (Dedyukhin 2012). **Republic of Tatarstan (TA)**: near Kazan in 1894 (Lebedev 1906). **Orenburg Oblast (ORE)**: vicinity of Orenburg in 1896 (Semenov-Tian-Shansky 1927).

The Asian part of the range, Russia (Fig. 3). **Sverdlovsk Oblast (SVE)**: Verkhnesaldinsky (1) and Novolyalinsky (2) districts. Conservation status: rare species (Gorbu-nov 2018). **Chelyabinsk Oblast (CHE)**: vicinity of Zlatoust in 1898 (additional material, ZIN). **Tyumen Oblast (TYU)**: Yarkovsky (1), Nizhnetavdinsky (2), Yalutorovsky (3), and Vagaysky (4) districts; vicinity of Tobolsk (5) (Kolosov 1933; Sitnikov 2000, 2004, 2013; Stolbov et al. 2019). Conservation status: rare species. **Khanty-Mansi Autonomous Okrug (KHM)**: Sovetsky District and Kondinskies Lakes Natural Park in 2006 (Ukhova 2009; Gashev et al. 2012). **Tomsk Oblast (TOM)**: Tomsky (1) and Verkhneketsky districts (2) (Kraatz 1879; Kiseleva 1926; Kuleshov and Romanenko 2009; Krivets and Vysotina 2011); Pervomaysky District (3) (additional material, ISEA). **Novosibirsk Oblast (NVS)**: Novosibirsk (1) (former Novonikolayevsk) in 1910; Bolshaya Elovka River (2) in 1912 (additional material, ZIN). **Kemerovo Oblast (KEM)**: Salair (1) and Novokuznetsk (2) (Gebler 1848); later found in Novokuznetsky District (3) (additional material, ISEA); 4 km north of the Sheregesh urban-type settlement Tashtagolsky District (4) (Efimov 2001, pers. comm.) where unpublished records were previously made (additional material, ISEA); Mezhdurechensky District (5) (Kazakova 2018). **Altai Krai (ALT)**: Tigirekskiy Nature Reserve (1) in 2014 (Gus'kova and Kuftina 2015); Troitsky District (2) (additional material, ISEA). Conservation status: very rare species (Gus'kova 2016). **Altai Republic (AL)**: near Lake Teletskoye (1) in 1901 (Semenov-Tian-Shansky 1927) and in 1969 (additional material, ISEA); Choysky (2) (additional material, ISEA), Shebalinsky (3) (additional material, ISEA), and Chemalsky (4) districts (additional material, ZIN); reported as common (Cherepanov 1979). **Republic of**

Khakassia (KK): Askizsky District (additional material, ISEA). **Krasnoyarsk Krai (KYA)**: Minusinsky District in the vicinity of Znamenka village (1) in 2009 (Listvyagova 2013); old record in the vicinity of Krasnoyarsk (2) (additional material, ZIN). **Irkutsk Oblast (IRK)**: Ziminsky (1), Tayshetsky (2), Usolsky (3), Shelekhovskiy (4), Slyudyansky (5), and Irkutsky (6) districts (Kraatz 1879; Agafonova and Antonov 2014). **Republic of Buryatia (BU)**: south of Lake Baikal in the Kaban-sky District in 1963 (Agafonova and Antonov 2014) and in 1973 (Švácha and Danilevsky 1986). **Zabaykalsky Krai (ZAB)**: Shilkinsky District (new record). **South of the Russian Far East: Amur Oblast (AMU), Jewish Autonomous Oblast (YEV), Khabarovsk Krai (KHA), and Primorsky Krai (PRI)**: reported to occur in this territory (Samoylov 1936; Shabliovskiy 1956; Bílý and Mehl 1989) but not confirmed.

Tragosoma depsarium, a boreomontane species, is predominantly found in the foothill and mountain sub-boreal and boreal forests of Eurasia and it sporadically distributed throughout its range.

In Europe, *T. depsarium* has been studied since its first description by Linné (1767). The Swedish population of *T. depsarium* is currently quite large and stable (Jeppsson et al. 2010). *Tragosoma depsarium* is also found relatively regularly found in the Alps, Pyrenees, Carpathians, and mountains of the Balkan Peninsula. Outside of these mountainous regions and Scandinavia, the largest number of new records was made in central and eastern European Russia. In of its European range, the most recent records were made over 20 years ago, and sometimes even more than 100 years ago. Today, *T. depsarium* definitely does not occur in Belgium, and its presence in Poland, Italy, Ukraine, and Estonia needs to be confirmed. The status of *T. depsarium* populations within Russia, in the Pskov, Bryansk, Kaluga, and Orenburg Oblasts and the Republic of Tatarstan is currently unknown; all published records from these territories were made a very long time ago. The presence of the species in the Caucasus (Bílý and Mehl 1989) is unconfirmed (Danilevsky 2014).

Much less is known about the distribution of *T. depsarium* in Siberia, as entomological studies here were started later and then on a smaller scale and in less detail than in Europe. The vastness of Siberia and the poor development of transport infrastructure is the cause. Most of the records are known from in the western part of the West Siberian Plain: Sverdlovsk, Tyumen, Tomsk and Kemerovo Oblasts and in Altai. We have found no published records of *T. depsarium* made in the lowlands of the central West Siberian Plain. It is unclear whether this is due to habitat discontinuity or insufficient investigations.

Many records of *T. depsarium* have been made near Lake Baikal, especially in the Irkutsk Oblast. The records of larvae south of Lake Baikal have for a long time been considered the easternmost ones (Danilevsky

2014). Many researchers reported that this species occurs east of Lake Baikal to the Pacific Ocean, along the Amur or the northern borders of China (Hamilton 1892; Saalas 1923; Plavilstshikov 1936; Samoylov 1936; Gressitt 1951; Bílý and Mehl 1989). Shabliovsky (1956) included the Khabarovsk Krai and Primorsky Krai in the range of *T. depsarium* and mentioned that the species develops on *Picea jezoensis* (Siebold & Zucc.) Carr., *Picea obovata* Ledeb., and *Pinus koraiensis* Siebold & Zucc. and provided detailed descriptions of the shape of larval galleries. However, none of these authors have reported records based on confirmed material from the Russian Far East. We failed to find specimens of *T. depsarium* in the entomological collection of the Federal Scientific Center of the East Asia Terrestrial Biodiversity FEB RAS. According to Danilevsky (2014), the occurrence of *T. depsarium* in the Russian Far East and in North China cannot be considered established. Today, the easternmost record of this species is that made in the Zabaykalsky Krai and presented in this study. Substantiation of the occurrence of *T. depsarium* in the Russian Far East requires additional research.

Hence, based on our study of reliable materials, *T. depsarium* has a western–central Palaearctic range, and with the greatest number of records in the west. This species' farther east in Siberia, near the Pacific Ocean, remains to be proved.

Acknowledgements

We are grateful for the help, advice, and valuable information from our colleagues from Serbia: I. Gnjatovic (University of Niš), F. Vukajlović (University of Kragujevac), D. Pavićević (Belgrade); Bulgaria: B. Gueorguiev (National Museum of Natural History, Sofia); Greece: G. Kakiopoulos (Athens); Hungary: O. Merkl (Hungarian Natural History Museum, Budapest), N. Rahmé (Budapest); Lithuania: V. Tamutis (Vytautas Magnus University, Kaunas); Latvia: D. Telnov (University of Latvia, Riga; Natural History Museum, London); Estonia: I. Süda (Estonian University of Life Sciences, Tartu); Ukraine: A.M. Zamoroka (Vasyl Stefanyk Precarpathian National University, Ivano-Frankivsk); Belarus: M.A. Lukasheva (Baranovich State University), A.V. Derunkov (SPC of the NAS of Belarus for Biological Resources); Russia: M.L. Danilevsky and A.O. Bienkowski (A.N. Severtsov Institute of Ecology and Evolution, Moscow), L.V. Egorov (State Nature Reserve Prisurevsky, Cheboksary), A.I. Miroshnikov (Sochi National Park, Sochi), A.S. Sazhnev (I.D. Papanin Institute for biology of inland waters, Borok), R.N. Ishin (Tambov), P.S. Sitnikov (Tyumen), D.A. Efimov (Kemerovo State University), N.L. Ukhova (Visim Nature Reserve, Kirovgrad), E.S. Koshkin (Institute of Water and Ecology Problems, Khabarovsk), and E.V. Novomodnyi (Khabarovsk Territorial Museum named after N.I. Grodekov).

We are also very grateful to M.G. Volkovitch (Zoological Institute of the Russian Academy of Sciences,

Saint Petersburg), S.E. Tshernyshev, A.A. Legalov and V.K. Zinchenko (Institute of Systematics and Ecology of Animals, Novosibirsk), A.S. Lelej and S.Yu. Storozhenko (Federal Scientific Center of the East Asia Terrestrial Biodiversity, Vladivostok) for providing an opportunity to study the collections of the mentioned scientific institutions. We would like to thank V.A. Golovizin (Krasnoyarsk) for the *T. depsarium* specimens provided for the study.

Authors' Contributions

VGB identified the species. NSA and VGB studied the material from the collections. NSA searched the literature, photographed the specimens collected from Zabaykalsky Krai, and prepared the distribution map. All authors have read and approved the final version of the manuscript. The authors declare that they have no conflict of interest.

References

- Adakhovsky DA, Dedyukhin SV (2011) Invertebrates. In: Baranova OG (Ed.) Rare and endangered plant and animal species in the southern half of Udmurtia and their protection: Results of scientific research (2005–2009). Publishing House “Udmurtskiy Universitet”, Izhevsk, Russia, 92–112. [in Russian]
- Adlbauer K (1992) Die Bockkäfer des Fürstentums Liechtenstein (Col., Cerambycidae). Bericht der Botanisch-Zoologischen Gesellschaft Liechtenstein-Sargans-Werdenberg 19: 253–293.
- Adlbauer K (2001) 2. Nachtrag zur Bockkäferfauna der Steiermark unter dem Aspekt der Artenbedrohung (Coleoptera, Cerambycidae). *Joannea Zoologie* 3: 83–104.
- Adlbauer K (2010) Die Bockkäfer des Nationalpark Gesäuse. *Coleoptera, Cerambycidae. Joannea Zoologie* 11: 51–95.
- Agafonova TA, Antonov IA (2014) Catalog of xylophagous insects of conifers of Baikal Siberia and Northern Amur Region: based on the stock materials of the Museum “Terrestrial Ecosystems of Baikal Siberia” SIPPB SB RAS. Institute of Geography SB RAS, Irkutsk, Russia, 215 pp. [in Russian]
- Alekseev SK, Perov VV (2017) *Tragosoma depsarium* (Linnaeus, 1767). In: Antochina VA et al. (Eds.) Red Data Book of the Kaluga Oblast. Volume 2. Animal world. Vash Dom, Kaluga, Russia, 93. [in Russian]
- Apfelbeck V (1894) Fauna insectorum balcanica. Beiträge zur Kenntnis der Balkanfauna. Wissenschaftliche Mitteilungen aus Bosnien und der Herzegowina 2: 511–542.
- Arnold NM (1902) A catalogue of the insects of Mogilev province. Ministry of Public Education, Sankt-Petersburg, Russia, 150 pp. [in Russian]
- Aurenhammer S, Komposch C, Schneider M, Degasperi G (2019) Urwaldrelikte Kärntens – Käfergemeinschaften von Naturwäldern im Spannungsfeld zwischen Forstwirtschaft und Naturschutz (Insecta: Coleoptera). *Carinthia* 2 (209/129): 431–466.
- Berger P, Kakiopoulos G, Brustel H, Minetti R (2010) Contribution à la connaissance des cerambycides (Coleoptera, Cerambycidae) de Grèce: 5eme note. *Biocosme Mesogéen* 27 (1): 17–26.
- Bertolini S (1895) Contribuzione alla fauna trentina dei coleotteri. *Bullettino della Società entomologica italiana* 27: 117–168.
- Bílý S, Mehl O (1989) Longhorn beetles (Coleoptera, Cerambycidae) of Fennoscandia and Denmark. *Fauna Entomologica Scandinavica* 22: 1–203.
- Binot M, Bless R, Boye P, Gruttke H, Pretschner P (Eds.) (1998) Rote Liste gefährdeter Tiere Deutschlands. Schriftenreihe für Land-

- schaftspflege und Naturschutz 55: 1–434 pp.
- Blanc M (2011) Redécouverte de *Tragosoma depsarium* (Linnaeus, 1767) dans les Alpes françaises (Coleoptera, Cerambycidae). *Entomologia Helvetica* 4: 85–87.
- Blindheim T (2020) BioFokus. Version 1.1592. BioFokus. Occurrence dataset <https://doi.org/10.15468/jxbhgx>. Accessed via GBIF.org on: 2020-12-09.
- Brandstetter CM, Kapp A (1998) Käferinventar von Vorarlberg und Liechtenstein, Insecta: Coleoptera. Eigenverlag des EVCV, Bürs, Austria, 92 pp.
- Breljih S, Drovenik B, Pirnat A (2006) Material for the Beetle Fauna (Coleoptera) of Slovenia. 2nd contribution: Polyphaga: Chrysomeloidea (= Phytophaga): Cerambycidae. *Scopolia* 58: 1–442.
- Brin A, Brustel H, Valladares L (2006) Contribution à la connaissance des coléoptères saproxyliques de la vallée du Marcadau (Hautes-Pyrénées). *Bulletin de la Société Linnéenne de Bordeaux* 140 (N.S.) 34 (1): 55–64.
- Brustel H, Freeman JC, Valladares L, Van Meer C (2001) Données originales sur quelques Cerambycidae des Pyrénées et régions voisines (Coleoptera). *Bulletin de la Société Linnéenne* 29 (1): 11–20.
- Cherepanov AI (1979) Longicorn beetles of North Asia (Prioninae, Desteniinae, Lepturiinae, Aseminae). *Nauka, Novosibirsk, Russia*, 472 pp. [in Russian]
- Cherkunov N (1889) List of beetles found in Kyiv and its environs. *Notes of the Kyiv Society of Naturalists* 10 (1): 147–204. [in Russian]
- Chernyshov AP (1930) List of beetles of the former Kaluga province. In: *Insect fauna of the former Kaluga province. Volume 2. Kaluga, Russia*, 5–16. [in Russian]
- Chistovsky SM (1929) List of Coleoptera (beetles) of Pskov and Velikie Luki districts. In: *Collected works of the Pskov Society of Local Lore. Volume 4. Pskov, Russia*, 89–112. [in Russian]
- Chittaro Y, Monnerat C, Sanchez A, Gonseth Y (2020) Swiss National Coleoptera Databank. Version 1.4. Swiss National Biodiversity Data and Information Centres – infospecies.ch. Occurrence dataset <https://doi.org/10.15468/cjmjy2>. Accessed via GBIF.org on: 2020-12-12.
- Dabasdati.lv (2020) Latvian citizen science portal. <https://dabasdati.lv/observation/cemlej7sea72g29abi4gud88pl/>. Accessed on: 2020-12-13.
- Danilevsky ML (2005) A checklist of the longicorn-beetles (Coleoptera: Cerambycidae) of Moscow Region. *Russian Entomological Journal* 15 (1): 43–51.
- Danilevsky ML (2014) Longicorn beetles (Coleoptera, Cerambycoidea) of Russia and adjacent countries. Part 1. HSC, Moscow, Russia, 522 pp. [in Russian with English and German abstract]
- Dedyukhin SV (2005) The longicorn beetles (Coleoptera, Cerambycidae) of National Park “Netchkinskiy” (with review of the family fauna of Udmurtia). *Vestnik of Udmurtia University* 10: 81–96. [in Russian with English abstract]
- Dedyukhin SV (2006) The data on the Red Data Book Coleoptera (Insecta, Coleoptera) and the species recommended to a new red list in the Udmurt Republic. *Vestnik of Udmurtia University* 10: 129–140. [in Russian with English abstract]
- Dedyukhin SV (2007) New data on the Udmurt fauna and spreading of longicorn beetles (Coleoptera: Cerambycidae). *Vestnik of Udmurtia University* 10: 65–69. [in Russian with English abstract]
- Dedyukhin SV (2012) *Tragosoma depsarium* (Linnaeus, 1767). In: Baranova OG (Ed.) *Red Book of the Udmurt Republic. Second edition. Perfektum, Cheboksary, Russia*, 50. [in Russian]
- Demelt CV (1957) Interessante Beobachtungen am Bockkäfer *Tragosoma depsarium* L. in Kärnten. *Carinthia* 2 (67): 139–143.
- Dolgin MM, Kolesnikova AA (2019) *Tragosoma depsarium* (Linnaeus, 1767). In: Degteva SV (Ed.) *Red Data Book of the Komi Republic. Third edition, official. Komi Republican Printing House, Syktyvkar, Russia*, 608. [in Russian]
- Drumont A, Komiya Z (2010) Cerambycidae: Prioninae. In: Löbl I, Smetana A (Eds.) *Catalogue of Palaearctic Coleoptera. Volume 6. Chrysomeloidea*. Apollo Books, Stenstrup, Denmark, 86–95.
- Duhem P, Horellou A (2020) Inventaire National du Patrimoine Naturel. Données photographiques validées de la galerie du forum “Le Monde des Insectes”. Version 1.1. UMS PatriNat (OFCN-CNRS-MNHN), Paris. Occurrence dataset. <https://doi.org/10.15468/2vjhg8>. Accessed via GBIF.org on: 2020-12-11.
- Eckelt A, Degasperi G (2018) Zur Diversität der xylobionten Käferfauna (Insecta: Coleoptera) des Samina- und Galinatales (Österreich und Liechtenstein) – Einblicke und Schlussfolgerungen. *Inatura Forschung online* 58: 1–20 pp.
- Efimov DA (2001) To the fauna of longicorn beetles (Coleoptera, Cerambycidae) of the Kemerovo Oblast. In: Bannih SV, Bibik OI, Elgina SI, Chernov SV, Shteinpreis TA (Eds.) *Proceedings of the regional scientific conference “Young scientists – for Kuzbass. A look into the XXI century”*. Kemerovo State Medical Academy, Kemerovo, Russia, 65–70. [in Russian]
- Egorov LV (2009) Materials on Coleoptera of the Red Data Book of the Chuvash Republic. In: Popov II, Kozlov VA, Samartsev VV, Zinov VG, Yashin VG (Eds.) *Materials of the Fourth International Scientific School “Science and Innovation - 2009” ISS “SI-2009”*: Materials of the Fourth International Scientific Seminar “Fundamental Research and Innovation” and the All-Russian Youth Scientific Seminar “Science and Innovation - 2009”. Mari State University, Yoshkar-Ola, Russia, 296–304. [in Russian]
- Egorov LV (2010) *Tragosoma depsarium* (Linnaeus, 1767). In: Isaev IV (Ed.) *Red Data Book of the Chuvash Republic. Volume 1. Part 2. Rare and endangered animal species. State Unitary Enterprise “IPK” Chuvashia*, Cheboksary, Russia, 47. [in Russian]
- Egorov LV (2018) New information on the beetle fauna (Insecta, Coleoptera) of Chuvashia. Report 9. In: Anikin VV, Gorshkov YuA, Egorov LV, Zolotukhin VV, Sinev SYu, Shulaev NV (Eds.) *Proceedings of the Kazan Branch of the Russian Entomological Society. Vol. 5. Materials of reports of the II Readings in memory of Professor Eduard Friedrich Eversmann*. Olitech, Kazan, Russia, 13–21. [in Russian]
- European Nucleotide Archive (EMBL-EBI) (2019) Geographically tagged INSDC sequences. Occurrence dataset. <https://doi.org/10.15468/cndomv>. Accessed via GBIF.org on: 2020-12-12. <https://www.gbif.org/occurrence/2311108126>
- Flossmann S (2006) Faunistisch interessante Funde von Bockkäfern (Coleoptera, Cerambycidae) in Deutschland. *Entomologische Nachrichten und Berichte* 50: 170–172.
- Foit J (2007) A species of longhorn beetle (Coleoptera: Cerambycidae) new to the Albanian fauna—new record of *Tragosoma depsarium* (L.). *Acta Entomologica Serbica* 12 (1): 87–89.
- Gashev SN, Zherebiatjeva NV, Korotkikh NN, Larin EG, Levashova MV, Popova TV, Stavishenko IV, Uvarova VI, Ukhova NL, Khoroshavin VYu, Khrenov VYa, Khritan’ko OA (2012) *Natural Park “Kondinskies Lakes”*. Ural Publishing Polygraphic Center, Yekaterinburg, Russia, 398 pp. [in Russian]
- GBIF Secretariat (2020). GBIF backbone taxonomy. <https://doi.org/10.15468/39omei>. Accessed via <https://www.gbif.org/species/5284517>. Accessed on: 2020-12-9.
- Gebler FA von (1848) Verzeichniss der im Kolywano-Woskresenskischen Hüttenbezirke Süd-west Sibiriens beobachteten Kaefer mit Bemerkungen und Beschreibungen. *Bulletin de la Société Impériale des Naturalistes de Moscou* 21 (2): 317–423.
- Ghiliani V (1887) Elenco delle specie di Coleotteri trovate in Piemonte (opera postuma a cura di L. Camerano). *Annali della Reale Accademia d’agricoltura di Torino* 29: 347–381.
- González C, Vives E, Zuzarte A (2007) Nuevo catálogo de los Cerambycidae (Coleoptera) de la Península Ibérica, islas Baleares e islas atlánticas: Canarias, Açores y Madeira. *Monografías Sociedad Entomológica Aragonesa* 12: 1–211.
- Gorbunov PYu (2018) *Tragosoma depsarium* (Linnaeus, 1767). In: Korytin NS (Ed.) *Red Data Book of the Sverdlovsk Oblast: animals, plants, mushrooms*. Mir, Yekaterinburg, Russia, 126. [in Russian]

- Gressitt JL (1951) Longicorn Beetles of China. In: Lepesme P (Ed.) Longicornia. Volume 2. Paul Lechevalier, Paris, France, 1–667.
- Guéorguiev B (2015) Gueorguiev's Coleoptera assessed. In: Golemansky V, Stoev P, Dobrev D, Beron P, Zivkov M, Popov A, Popov V, Beschkov V, Deltchev C, Michev T, Spassov N (Eds.) Red Data Book of the Republic of Bulgaria. Volume 2. Animals. BAS & MOEW, Sofia, Bulgaria, 353–358.
- Gus'kova EV, Kuftina GN (2015) Record of the rare longhorn beetle *Tragosoma depsarium* (Linnaeus, 1767) (Coleoptera, Cerambycidae) in the Tigireksky strict nature reserve (Altai Krai). Amurian Zoological Journal 7 (2): 134–136. [in Russian with English summary]
- Gus'kova EV (2016) *Tragosoma depsarium* (Linnaeus, 1767). In: Irisova NL, Shapetko EV, Yakovlev RV (Eds.) Red Data Book of Altai Krai. Volume 2. Rare and endangered species of animals. Altai University Publishing House, Barnaul, Russia, 42–43. [in Russian]
- Gutowski JM (2004) *Tragosoma depsarium* (Linnaeus, 1767). Grac. In: Głowaciński Z, Nowacki J (Eds.) Polska Czerwona Księga Zwierząt. Bezkręgowce. Instytut Ochrony Przyrody PAN, Akademia Rolnicza im. Cieszkowskiego, Krakow/Poznan, Poland, 142–143.
- Hamilton J (1892) Notes on Coleoptera—no. XI. The Canadian Entomologist 24 (12): 293–300.
- Hellrigl KG (1967) Die Cerambyciden-Fauna von Südtirol. Koleopterologische Rundschau 5 (45): 3–75.
- Hellrigl KG (2010) Faunistik der Bockkäfer von Südtirol (Coleoptera: Cerambycidae). Forest Observer 5: 31–152.
- Holecová M, Franc V (2001) Červený (ekozozologický) zoznam chrobákov (Coleoptera) Slovenska. In: Baláž D, Marhold K, Urban P (Eds.) Red List of plants and animals of Slovakia Nature Conservation. Ochrana prírody 20 (suppl.). Štátna Ochrana Prírody Slovenskej Republiky, Banská Bystrica, Slovakia, 111–128.
- Humala AE, Polevoi AV (2009) On the insects fauna of South-East Karelia. Transactions of Karelian Research Centre of Russian Academy of Science 9 (4): 53–75. [in Russian with English abstract]
- Ilić N, Čurčić S (2015) A checklist of longhorn beetles (Coleoptera: Cerambycidae) of Serbia. Zootaxa 4026 (1): 1–97. <https://doi.org/10.11646/zootaxa.4026.1.1>
- Ivshin NV (1990) Species composition and distribution of coleopteran insects of the Cerambycidae family on the territory of the Mari ASSR. In: Matveev VA (Ed.) Fauna and ecology of animals of the Middle Volga region: interuniversity collection of scientific papers. Margosuniversitet, Yoshkar-Ola, Russia, 9–16. [in Russian]
- Jäch MA (1994) Rote Liste der gefährdeten Käfer Österreichs (Coleoptera). In: Gepp J (Ed.) Rote Listen gefährdeter Tiere Österreichs. Grüne Reihe des Bundesministeriums für Umwelt, Jugend und Familie, Band 2. Graz, Austria, 107–200.
- Jakovlev JB (2007) *Tragosoma depsarium* (Linnaeus, 1767). In: Ivanter EV, Kuznetsov OL (Eds.) Red Data Book of Republic Karelia. Karelia, Petrozavodsk, Russia, 282–283. [in Russian]
- Jakovlev JB, Humala AE, Polevoi AV (2003) Insects. In: Gromtsev AN, Kitaev SP, Krutov VI, Kuznetsov OL, Lindkholm T, Jakovlev JB (Eds.) Biotic diversity of Karelia: conditions of forming, associations, species. Karelian Scientific Center RAS, Petrozavodsk, Russia, 159–168. [in Russian]
- Jeppsson T, Lindhe A, Gärdenfors U, Forslund P (2010) The use of historical collections to estimate population trends: a case study using Swedish longhorn beetles (Coleoptera: Cerambycidae). Biological Conservation 143 (9): 1940–1950. <https://doi.org/10.1016/j.biocon.2010.04.015>
- Kabátek P, Skořepa L (2017) Cerambycidae (tesaříkovití). In: Hejda R, Farkač J, Chobot K (Eds.) Červený seznam ohrozených druhů České republiky. Bezobratlí. List of threatened species in the Czech Republic. Invertebrates. Příroda, volume 36. Nature Conservation Agency of the CR, Prague, Czech Republic, 302–304.
- Karjalainen S, Salokannel J, Albrecht A (2010) Atlas of the Beetles of Finland. Cerambycidae – Curculionidae. <http://koivu.luomus.fi/elaintiede/kovakuoriatset/CATLAS4.HTM>. Last accessed on: 2020-12-09.
- Kazakova S (2018) Species diversity of land beetles of the Kuznetsk Alatau in the Podnebesnye Zub'ya area. In: Materials of the scientific-practical conference of research works of students of educational institutions of the Kemerovo region “Erudit-2018”, Kemerovo, Russia, 278–280. [in Russian]
- Kiseleva EF (1926) About longicorn beetles (Coleoptera, Cerambycidae) in the vicinity of Tomsk. Izvestiya Tomskogo Gosudarstvennogo Universiteta 77 (2): 123–133. [in Russian]
- Kolosov YuM (1933) Materials for the insect fauna of the Tobolsk North. Proceedings of the Ural Regional Institute Microbiology and Epidemiology 1 (1): 49–58. [in Russian]
- Kraatz G (1879) Ueber die Bockkäfer Ost-Sibiriens, namentlich die von Christoph am Amur gesammelten. Deutsche entomologische Zeitschrift 23 (1): 77–117.
- Kráska A (2015) Ochrana saproxylického hmyzu a opatření na jeho podporu: metodika AOPK ČR. Nature Conservation Agency of the CR, Prague, Czech Republic, 156 pp.
- Krivets SA, Vysotina SV (2011) Ecological and faunistic review of the coleopterous (Insecta, Coleoptera) dendrophagans of the Priketye conifer forests (Tomsk oblast). Tomsk State University Journal of Biology 3 (15): 111–126. [in Russian with English abstract]
- Krivokhatsky VA (2002) *Tragosoma depsarium* (Linnaeus, 1767). In: Noskov GA (Ed.) Red Data Book of Nature of the Leningrad Region. Volume 3. Animals. World and Family, Saint Petersburg, Russia, 159–160.
- Kuleshov DA, Romanenko VN (2009) Longicorn beetles (Coleoptera, Cerambycidae) of the Tomsk region. Tomsk State University Journal of Biology 4 (8): 29–40. [in Russian with English abstract]
- Kust T (2016) Entomofaunistische Untersuchungen im Wildnisgebiet Dürrenstein. 1. Teil: Käfer (Coleoptera), Schwebfliegen (Diptera: Syrphidae), Köcherfliegen (Trichoptera). Silva Fera 5: 78–94.
- Lapenu P (2020) Inventaire National du Patrimoine Naturel. Observations occasionnelles Parc national des Pyrénées. Version 1.1. UMS PatriNat (OFB-CNRS-MNHN), Paris. Occurrence dataset. <https://doi.org/10.15468/g3k5au>. Accessed via GBIF.org on 2020-12-11.
- Laplante S (2017) Description of a new Nearctic species of *Tragosoma* Audinet-Serville (Coleoptera: Cerambycidae: Prioninae), with species validations, new synonymies and a lectotype designation. Insecta Mundi 0578: 1–17.
- Lebedev AG (1906) Materials for the fauna of beetles of Kazan gubernia. In: Semenov AP, Kuznetsov NYa (Eds.) Proceedings of the Russian Entomological Society. Volume 37. Saint Petersburg, Russia, 352–438. [in Russian]
- Linné C (1767) Systema naturae. Tom. I. Pars II. Editio duodecima reformata. Laurent Salvii, Stockholm, Sweden, 533–1327.
- Listvyagova NA, Raykhert AI, Skribchenko AV (2013) Catalog of the collection of longicorn beetles (Insecta, Cerambycidae, Lepyturini) of the zoological museum of the Khakass State University named after N.F. Katanova (report 2). Bulletin of Katanov Khakass State University 6: 23–29. [in Russian]
- Ljungberg H (2015) Rödlista över skalbaggar Coleoptera. In: Westling A (Ed.) Rödlistade arter i Sverige 2015. Artdatabanken, Uppsala, Sweden, 119–140.
- Ljungberg H, Andrén B, Ehnström B, Jansson N, Jonsell M, Lindelöw Å, Lundberg S, Malmqvist A, Nilsson SG, Pettersson RB, Ramqvist T, Wanntorp HE, Wikars LO (2010) Skalbaggar – beetles coleoptera. In: Gärdenfors U (Ed.) The 2010 Red List of Swedish species. Artdatabanken, Uppsala, Sweden, 411–451.
- Lovcik P (1970) Bisherige Bockkäferfunde (Cerambycidae) aus dem Lande Salzburg. In: Mitteilungen aus dem Haus der Natur Salzburg. Festschrift der Naturwissenschaftlichen Arbeitsgemeinschaft am Haus der Natur in Salzburg zum 80. Geburtstag von Prof. Dr. Eduard Paul Tratz, Salzburg, Austria, 52–72.
- Malmberg S, Mannerkoski I, Martikainen P, Clayhills T, Helve E, Hyvärinen E, Karjalainen S, Mattilaaal J, Muona J, Rassi P (2019)

- Beetles. In: Hyvärinen E, Juslén A, Kemppainen E, Uddström A, Liukko UM (Eds.) The 2019 Red List of Finnish species. Ministry of the Environment & Finnish Environment Institute, Helsinki, Finland, 391–425.
- Mannerkoski I, Hyvärinen E, Campanaro A, Alexander K, Büche B, Dodelin B, Mason F, Pettersson R, Mico E, Méndez M (2010) *Tragosoma depsarium*. The IUCN Red List of threatened species 2010: e.T157922A5174376. <https://www.iucnredlist.org/species/157922/5174376>. Accessed on: 2020-12-13.
- Martinelli A (1996) I cerambicidi della Val di Genova. *Annali del Museo Civico di Rovereto* 11: 349–360.
- Medvedev AA (1997) Fauna of longicorn beetles (Coleoptera, Cerambycidae) in the vicinity of the biological station of Syktyvkar State University. In: Oischwang VN, Bogacheva IA, Nikolaeva NV, Mikbailov YuE, Gorbunov PYu, Zinovjev EV (Eds.) Achievements of entomology in the Urals. Collection of scientific papers. Ekaterinburg, Russia, 186. [in Russian]
- Medvegy M (2001) Endangerment of the Hungarian longhorn beetles. *Természetvédelmi Közlemények* 9: 163–199.
- Mesaroš G (2020) Podaci o rasprostranjenju vrste *Tragosoma depsarium* (Linnaeus, 1767) u Srbiji. Portal za kartiranje biološke raznovrsnosti Srbije – BioRas. <http://bioras.petnica.rs/rasprostranjenost.php?id=1083>. Accessed on: 2020-12-13.
- Migliaccio E, Georgiev G, Gashtarov V (2007) An annotated list of Bulgarian Cerambycids with special view on the rarest species and endemics (Coleoptera: Cerambycidae). *Lambillionea* 107 (1): 3–79.
- Mikšić R (1963) Beitrag zur Kenntnis der Bockkäferfauna (Cerambycidae) Jugoslawiens. *Acta Biologica* 3: 55–188.
- Mikšić R, Georgijević E (1971) Cerambycidae of Yugoslavia. Part I. ANUBiH, Sarajevo, Bosnia and Herzegovina, 175 pp.
- Mitter H (1988) Besonderheiten aus der Käferfauna des Mühlviertels. In: Litschel H (Ed.) Das Mühlviertel: Natur, Kultur, Leben. Oberösterreichische Landesausstellung 1988. Land Oberösterreich, Amt der Oö. Landesregierung, Abt. Kultur, Linz, Austria, 219–226.
- Mokrousov MV (2008) Proposals for the introduction of some species of insects in the Red Data Book of the Nizhny Novgorod Region. In: Anufriev GA, Bakka AI, Bakka SV, Karyakin IV, Kiseleva NYu, Kurochkin DV (Eds.) Rare species of living organisms in the Nizhny Novgorod region: collection of working materials of the Commission for the Red Data Book of the Nizhny Novgorod Oblast. Issue 1. Nizhny Novgorod, Russia, 25–31. [in Russian]
- Monnerat C, Barbalat S, Lachat T, Gonseth Y (2016) Rote Liste der Prachtkäfer, Bockkäfer, Rosenkäfer und Schröter. Gefährdete Arten der Schweiz. Reihe Umweltvollzug No. 1622. Bundesamt für Umwelt, Bern, Switzerland, 118 pp.
- Moscardini C (1956) I Cerambicidi della Val di Genova (Trentino). *Studi Trentini di Scienze Naturali* 5 (33): 54–74.
- Müller-Kroehling S, Walentowski H, Kölling C (2009) Natürliche Fichtenwälder im Klimawandel – Hochgradig gefährdete Ökosysteme. *LWF Wissen* 63: 70–85.
- Nieto A, Alexander KNA (2010) European Red List of Saproxylid Beetles. Publications Office of the European Union, Luxembourg, 45 pp.
- Nikitsky NB (2008) *Tragosoma depsarium* (Linnaeus, 1767). In: Varlygina TI, Zubakin VA, Sobolev NA (Eds.) Red Data Book of the Moscow Oblast. Second edition, revised and enlarged. KMK Scientific Press, Moscow, Russia, 209. [in Russian]
- Nikitsky NB (2009) New and interesting finds of xylophilous and some other species of beetles (Coleoptera) in the Moscow Region and in Moscow. *Bulletin of the Moscow Society of Naturalists* 114 (5): 50–58. [in Russian with English summary]
- Nowicki M (1864) Przyczynek do owadniczej fauny Galicyi. Krakow, Poland, 87 pp.
- Ødegaard F, Vang R (2020) NINA insect database. Version 1.75. Norwegian Institute for Nature Research. Occurrence dataset. <https://doi.org/10.15468/cf4c2s>. Accessed via GBIF.org on: 2020-12-09.
- <https://www.gbif.org/occurrence/1341824765>
- Ødegaard F, Hanssen O, Olberg S (2015) Coleoptera. In: Henriksen S, Hilmo O (Eds.) Norsk rødliste for arter 2015. Artsdatabanken, Trondheim, 19–41. [https://artsdatabanken.no/Files/13973/Norsk_r_dliste_for_arter_2015_\(PDF\)](https://artsdatabanken.no/Files/13973/Norsk_r_dliste_for_arter_2015_(PDF)). Accessed on: 2020-11-15.
- Patkó F (2008) A Hargita-hegységi xilofág (fát fogyasztó) bogarak (Insecta Coleoptera). In: Muranyi J (Ed.) A Csiki Székely Múzeum Évkönyve 2007–2008. Pro-Print Kiadó, Miercurea Ciuc (Csíkszereda), Romania, 265–296.
- Pawłowski J, Kubisz D, Mazur M (2002) Coleoptera. In: Głowaciński Z (Ed.) Red list of threatened animals in Poland. Polish Academy of Sciences, Institute of Nature Conservation, Krakow, Poland, 88–110.
- Pileckis S, Monsevičius V (1997) Lietuvos fauna. Vabalai. T. 2. Mokslas, Vilnius, Lithuania, 216 pp.
- Plaige V (2019) Inventaire National du Patrimoine Naturel. Données du parc national de la Vanoise. Version 1.1. UMS PatriNat (OFB-CNRS-MNHN), Paris. Occurrence dataset. <https://doi.org/10.15468/4xudk8>. Accessed via GBIF.org on: 2020-12-11.
- Pavilstshikov NN (1936) Longhorn beetles. Part 1. In: Stackelberg AA (Ed.) Fauna of the USSR. Coleoptera. T. 21. Academy of Sciences of the USSR, Moscow/Leningrad, Russia, 612 pp. [in Russian]
- Potani DV (2014) *Tragosoma depsarium* L. In: Anufriev GA, Bakka SV, Kiseleva NYu (Eds.) Red Data Book of the Nizhny Novgorod Oblast. Volume 1. Animals. Second edition, revised and enlarged. DEKOM, Nizhny Novgorod, Russia, 252–253. [in Russian]
- Radkevich AI (1936) Material for the study of the BSSR entomofauna. Fauna of beetles of the northeastern part of Belarus (Coleoptera). In: Proceedings of the Vitebsk Pedagogical Institute. Part 1. Vitebsk, Belarus, 115–160. [in Russian]
- Rapuzzi P, Sama G (2012) Contributo alla conoscenza dei Cerambycidae di Albania (Coleoptera, Cerambycidae). *Atti del Museo Civico di Storia Naturale di Trieste* 55: 181–234.
- Rapuzzi P, Biscaccianti AB, Baviera C (2014) Famiglia Cerambycidae. In: Audisio P, Baviera C, Carpaneto GM, Biscaccianti AB, Battistoni A, Teofili C, Rondinini C (Eds.) Lista Rossa IUCN dei coleotteri saproxilici Italiani. Comitato Italiano IUCN e Ministero dell'Ambiente e della Tutela del Territorio e del Mare, Roma, Italy, 132 pp.
- Saalas U (1923) Die Fichtenkäfer Finnlands I–II. *Annales Academiae Scientiarum Fennicae* 22 (1): 349–432.
- Samoylov TP (1936) Materials on the bioecology of longicorn beetles of the South Ussuriysk Territory. In: Poretsky AS (Ed.) Proceedings of the Mountain–Taiga Station of the Far Eastern Branch of the USSR Academy of Sciences. Volume 1. Dal'giz, Khabarovsk, Russia, 217–237. [in Russian]
- Schillhammer H (1995) Bemerkenswerte Käferfunde aus Österreich (IV) (Coleoptera). *Koleopterologische Rundschau* 65: 229–232.
- Schmidl J, Bussler H (2003) Rote Liste gefährdeter Bockkäfer (Coleoptera: Cerambycidae) Bayerns. In: Rote Liste gefährdeter Tiere Bayerns. LfU series of publications 166, Munich, Germany, 150–153.
- Sedykh KF (1974) Fauna of the Komi ASSR. Invertebrates. Komi Book Publishing House, Syktyvkar, Russia, 192 pp. [in Russian]
- Semenov-Tian-Shansky AP (1927) *Analecta Coleopterologica*. XX. *Revue Russe d'Entomologie* 21 (3–4): 230–239.
- Shabliovsky VV (1956) Keys to the larvae of the main species of longicorn beetles—pests of the forests of the Far East. In: Proceedings of the Far Eastern Branch of the USSR Academy of Sciences. Zoological series. Volume 3. Dal'giz, Vladivostok, Russia, 111–135. [in Russian]
- Shapovalov AM (2018) *Tragosoma depsarium* (Linnaeus, 1767). In: Bublischenko YuN, Golubkov SM, Kijashko PV (Eds.) Red Data Book of the Leningrad Oblast. Animals. Papyrus, Saint Petersburg, Russia, 325–326. [in Russian]
- Sitonen J, Martikainen P, Kaila L, Mannerkoski I, Rassi P, Rütanen I (1996) New faunistic records of saproxylid Coleoptera, Diptera,

- Heteroptera, Homoptera and Lepidoptera from the Republic of Karelia, Russia. *Entomologica Fennica* 7: 69–76.
- Sitnikov PS (2000) Rare finds of longicorn beetles from Tyumen collections. In: Petrova VP (Ed.) Tyumen land: yearbook of the Tyumen regional museum of local lore, Tyumen, Russia, 216–224. [in Russian]
- Sitnikov PS (2004) *Tragosoma deparium* (Linnaeus, 1767). In: Petrova OA (Ed.) Red Data Book of the Tyumen Region: animals, plants, mushrooms. Ural University Publishing House, Yekaterinburg, Russia, 116. [in Russian]
- Sitnikov PS (2013) New finds of rare insects from the Red Data Book of the Tyumen Oblast. In: Gashev SN (Ed.) Materials to the second edition of the Red Data Book of the Tyumen Region. Tyumen-NIIgiprogaz, Tyumen, Russia, 95–100. [in Russian]
- Sláma M (1998) Tesaříkoviti – Cerambycidae České republiky a Slovenské republiky (Brouci – Coleoptera). Milan Slama, Krhanice, Czech Republic, 383 pp.
- Spuris Z (Ed.) (1998) Red Data Book of Latvia. Rare and threatened species of plants and animals. Vol. 4, Invertebrates. Institute of Biology, Riga, Latvia, 388 pp.
- Stark NK (1926) List of beetles of the Bryansk forest area. In: Proceedings of the Smolensk Society of Naturalists and Physicians at Smolensk State University. Volume 1. Smolensk, Russia, 77–92. [in Russian]
- Stolbov VA, Sergeeva EV, Lomakin DE, Sheykin SD (2019) A checklist of longicorn beetles (Coleoptera: Cerambycidae) of Tyumenskaya Oblast of Russia. *Eurasian Entomological Journal* 18 (3): 199–212. <https://doi.org/10.15298/euroasentj.18.3.10>.
- Süda I (2008) Red Data Book of Estonia. Commission for Nature Conservation of the Estonian Academy of Sciences. <http://vana.elurikkus.ut.ee/prmt.php?lang=eng>. Accessed on: 2020-12-13.
- Švácha P, Danilevsky ML (1986) Cerambycoid Larvae of Europe and Soviet Union (Coleoptera, Cerambycoidea). Part I. *Acta Universitatis Carolinae – Biologica* 30: 1–176.
- Svetlov SA (1996) Fauna and ecology of the longicorn beetles of the Ivanovo Oblast. Graduate work. Ivanovo State University, Ivanovo, Russia, 72 pp. [in Russian]
- Telnov D, Kalniņš M (2003) To the knowledge of Latvian Coleoptera. 3. *Latvijas Entomologs*, 40: 21–33.
- Telnov D, Vilks K, Piterāns U, Kalniņš M, Fägerström C (2011) Contributions to the knowledge of Latvian Coleoptera 9. *Latvijas Entomologs* 50: 20–26.
- Thomson CG (1866) Skandinaviens Coleoptera, synoptiskt bearbetade. T. 8. Tryckt Ut Lundbergska Boktryckeriet, Lund, Sweden, 409 pp.
- Tikhomirov AM (2007) *Tragosoma deparium* (Linnaeus, 1767). In: Isaev VA (Ed.) Red Data Book of the Ivanovo Oblast. Volume 1. Animals. PresSto, Ivanovo, Russia, 40. [in Russian]
- Touroult J (2020) Inventaire National du Patrimoine Naturel. Collection entomologique de Patrick Jaumes (coléoptères). UMS PatriNat (OFB-CNRS-MNHN), Paris. Occurrence dataset. <https://doi.org/10.15468/3phjta>. Accessed via GBIF.org on: 2020-12-11.
- Tsinkevich VA (2015) *Tragosoma deparium* (Linnaeus, 1767). In: Kachanovsky IM, Nikiforov ME, Parfenov VI (Eds.) Red Data Book of the Republic of Belarus. Animals: rare and endangered species of wild Animals. Petrus Brovka Belarusian Encyclopedia Publishing House, Minsk, Belarus, 197. [in Russian]
- Turpaud A (2018) Inventaire National du Patrimoine Naturel. Données naturalistes d'Anthony Turpaud. UMS PatriNat (OFB-CNRS-MNHN), Paris. Occurrence dataset. <https://doi.org/10.15468/rsmj>. Accessed via GBIF.org on: 2020-04-01.
- Ukhova NL (2009) Specially Protected Species of Invertebrates in the Natural Park “Kondinskies Lakes”. In: Fedorov MYu (Ed.) Protection, environmental education, recreational activities, study of biodiversity and cultural heritage in protected areas: Materials of the regional scientific and practical conference dedicated to the 5th anniversary of the Natural Park “Chusovaya River”, Nizhny Tagil, Russia, 132–136. [in Russian]
- Vershelde D, Brosens D (2019a) Ghent University – Zoology Museum – Insect collection. Ghent University Occurrence dataset <https://doi.org/10.15468/6fcelj>. Accessed via GBIF.org on: 2020-12-12. <https://www.gbif.org/occurrence/1315922626>
- Vershelde D, Brosens D (2019b) Ghent University – Zoology Museum – Insect collection. Ghent University. Occurrence dataset. <https://doi.org/10.15468/6fcelj>. Accessed via GBIF.org on: 2020-12-12. <https://www.gbif.org/occurrence/1315922599>
- Vilks K, Pilāts M (2008) Notes on recent observation of rare longhorn beetle *Tragosoma deparium* (Linnaeus, 1767) (Coleoptera, Cerambycidae) in the Slītere National Park, Latvia. *Latvijas Entomologs* 46: 76–78.
- Vives E (1984) Cerambycoides (Coleoptera) de la Península Ibérica y de las Islas Baleares. *Treballs del Museu de Zoologia de Barcelona* 2. Museu de Zoologia, Barcelona, Spain, 137 pp.
- Vlasov DV (2015) *Tragosoma deparium* (Linnaeus, 1767). In: Nyankovsky MA (Ed.) Red Data Book of the Yaroslavl Oblast. Academy 76, Yaroslavl, Russia, 252–253. [in Russian]
- Vlasov DV (2019) The longhorn beetles (Coleoptera: Cerambycidae) of Yaroslavl Region. In: Egorov LV (Ed.) Scientific proceedings of the State Nature Reserve “Prisursky”. Volume 34. Cheboksary, Russia, 86–114. [in Russian with English summary]
- Voolma K, Süda I (1999) Rare species of Cerambycidae and Scolytidae (Coleoptera) in Estonia. *Proceedings of the XXIV Nordic Congress of Entomology*, Tartu, Estonia, 187–192.
- Wikars LO (2003) *Tragosoma deparium*. Coleoptera: Cerambycidae is temporarily favoured by clear-cuts but depends on old-growth forest. *Entomologisk Tidskrift* 124: 1–12.
- Wikars LO (2004) Habitat requirements of the pine wood-living beetle *Tragosoma deparium* (Coleoptera: Cerambycidae) at log, stand, and landscape scale. *Ecological Bulletins* 51: 287–294.
- Yuferev GI (2014) *Tragosoma deparium* (Linnaeus, 1767). In: Baranova OG, Lachokha EP, Ryabov VM, Sotnikov VN, Tarasova EM, Tselishcheva LG (Eds.) Red Data Book of the Kirov Oblast: animals, plants, mushrooms. Second edition. Kirov Regional Printing House, Kirov, Russia, 107. [in Russian]
- Yurkina EV (2011) Xylophagous insects – inhabitants of forests, timber and lumber of middle taiga forests of the Komi Republic. *Forestry Bulletin* 4 (80): 24–34. [in Russian]
- Zagaykevich IK (1986) Longhorn beetles (Coleoptera, Cerambycidae) of Belovezhskaya Pushcha. In: Zagaykevich IK, Kaziuchits AV (Eds.) Problems of protection of the gene pool and management of ecosystems in the reserves of the forest zone. Abstracts of the All-Union Council, Berezinsky Reserve. Part 2. Moscow, Russia, 86–87. [in Russian]
- Zamoroka AM (2018) The longhorn beetles (Coleoptera: Cerambycidae) of the Eastern Carpathian Mountains in Ukraine. *Munis Entomology & Zoology* 13 (2): 655–691.
- Zawadzki Z (1937) Kózki ziemi Wileńskie. *Spostrzeżenia i notaty. Polskie Pismo Entomologiczne* 14–15: 281–305.