

ISSN 0749-8004

JOURNAL OF  
**ENTOMOLOGICAL**  
S • C • I • E • N • C • E

**Distribution of *Rhaesus serricollis* (Coleoptera: Cerambycidae: Prioninae) in the Mediterranean Region<sup>1</sup>**

Sakine Serap Avgın<sup>2</sup>, Arvīds Barševskis<sup>3</sup>, and Uldis Valainis<sup>3</sup>

Kahramanmaraş Sütçü İmam University, Faculty of Education, Division of Science Education, and Faculty of Art and Science, Department of Biology, Avşar Campus, 46100 Kahramanmaraş, Turkey

# Distribution of *Rhaesus serricollis* (Coleoptera: Cerambycidae: Prioninae) in the Mediterranean Region<sup>1</sup>

Sakine Serap Avgın<sup>2</sup>, Arvīds Barševskis<sup>3</sup>, and Uldis Valainis<sup>3</sup>

Kahramanmaraş Sütçü İmam University, Faculty of Education, Division of Science Education, and Faculty of Art and Science, Department of Biology, Avşar Campus, 46100 Kahramanmaraş, Turkey

---

J. Entomol. Sci. 50(3): 206–217 (July 2015)

**Abstract** *Rhaesus serricollis* (Motschulsky) (Coleoptera: Cerambycidae: Prioninae), a near-threatened species on the European Red List of Saproxyllic Beetles, is reported from Dalmatia, Albania, Serbia, Macedonia in the Balkan peninsula to southern Greece, Bulgaria, Turkey, Iran, Syria, Israel, Cyprus, Caucasus, Georgia, Transcaucasia, the Near East, Lebanon, and North Africa. It also has been introduced into Egypt. In Turkey, it occurs in Adana, Antalya, Bilecik, Burdur, Denizli, Düzce, Hatay, İçel, İstanbul, İzmir, Kahramanmaraş, Konya, Muğla, and Osmaniye provinces and the Toros Mountains. Known host plants include *Castanea*, *Celtis* (Cannabaceae), *Fagus* (Fagaceae), *Ficus* (Moraceae), *Liquidambar* (Altingiaceae), *Liquidambar orientalis*, *Juglans* (Juglandoideae), *Juglans regia*, *Morus* (Moraceae), *Morus alba*, *Pinus brutia* (Pinaceae), *Pinus nigra*, *Platanus* (Platanaceae), *Platanus orientalis*, *Populus alba* (Salicaceae), *Quercus* (Fagaceae), *Quercus calliprinos*, *Quercus ithaburensis*, *Salix* (Salicaceae), and *Tilia* (Malvaceae).

**Key Words** *Rhaesus serricollis*, long-horned beetle, distribution, host plants

---


*Rhaesus serricollis* (Motschulsky) (Coleoptera: Cerambycidae: Prioninae) is classified as “near threatened” on the European Red List of Saproxyllic Beetles (Nieto and Alexander 2010). Assimilation of information on this species is critical to efforts to protect this species from extinction in the Mediterranean basin. The Mediterranean basin extends approximately 3,800 km east to west from the tip of Portugal to the shores of Lebanon and approximately 1,000 km north to south from Italy to Morocco and Libya. Within the European Union, the Mediterranean Region encompasses seven countries, either partially (France, Portugal, Italy, Spain) or completely (Greece, Malta, Cyprus) (Sundseth and Brussels 2009). The region is recognized for its floral and faunal biodiversity. Between 15,000 and 25,000 species of plants are distributed there, 60% of which are unique to the region. About one-third of the Mediterranean fauna is endemic (International Union for Conservation of Nature 2012). In many cases, information is incomplete regarding trends in biodiversity, especially with marine organisms, in the region; yet, there are sufficient data demonstrating that biodiversity in the region is at severe risk. According to the International Union for Conservation of Nature Red List of Threatened Species,

---

<sup>1</sup>Received 20 November 2014; accepted for publication 16 December 2014.

<sup>2</sup>Corresponding author (email: serapavgin@hotmail.com).

<sup>3</sup>Daugavpils University, Institute of Systematic Biology, Vienības Str., 13, Daugavpils, LV-5401, Latvia.



**Fig. 1. Distribution of *Rhaesus serricollis* in provinces of Turkey, Mediterranean basin, and other countries.**

19% of faunal species (amphibians, birds, cartilaginous fishes, endemic freshwater fishes, crabs and crayfish, mammals, dragonflies, and reptiles) are threatened with extinction (5% Critically Endangered, 7% Endangered, 7% Vulnerable) (Vlachogianni et al. 2012). Insect diversity in the region is also high, with most representatives in the Order Coleoptera. The coleopteran fauna of Turkey alone is estimated to include >25,000 species (Koçak and Kemal 2009).

Turkey plays an important geographic and ecological role in safeguarding biodiversity within its borders as well as in neighboring countries and regions. Turkey is positioned at an intersection of geographical regions with large climatic and geographical gradients as well as a diversity of ecosystems and habitats (International Union for Conservation of Nature 2012, Kahraman et al. 2011). Biodiversity in Turkey is under severe pressure, and region-wide conservation efforts must be implemented to avoid extinction of threatened species. Such conservation efforts can be aided by identification of emblematic umbrella species to help market, advertise, and generate public support.

Of the many habitats and species threatened in Turkey, the deadwood habitat of native broadleaf tree species is one that is under severe pressure (Avcı et al. 2010, Coşkun et al. 2010, Gürkan et al. 2010). *Rhaesus serricollis* is among the saproxylic beetles of that habitat and, in Europe, is considered Near Threatened (Nieto and Alexander 2010). Thus, the primary objective of this paper is to define the known distribution of this insect along with information on its ecological habits and host plants. In addition, a bibliography of previous studies related to this species is included.

**Table 1. Distribution of *Rhaesus serricollis* in Turkey by province, with bibliographic citation of occurrence.**

Province	Locality	Citations
Adana	Unreported	Özdikmen 2008, Özdikmen et al. 2012
Adana	Bacalı–Karataş	Tozlu et al. 2002
Adana	Karaisahlı, Yedigözüleğen	Özdikmen 2011, Cihan 2013, Cihan et al. 2013
Antalya	Alanya	Demelt 1963, Lodos 1998, Özdikmen 2008, Özdikmen et al. 2012, Erdem and Çanakçıoğlu 1977, Çanakçıoğlu 1983, Çanakçıoğlu and Mol 1998, Svacha and Danilevsky 1987
Antalya	Alanya–Bambus Camp	Aldbauer 1988
Antalya	Central–Çaltıcak–Finike (Turunçova)–Serik	Tozlu et al. 2002
Antalya	Manavgat, Demirciler village	Özdikmen 2007
Antalya	Toros Mountains (Elmah)	Villiers 1967
Bilecik	Unreported	Bodemeyer 1906, Erdem and Çanakçıoğlu 1977, Çanakçıoğlu 1983, Çanakçıoğlu and Mol 1998
Burdur	Central	Tozlu et al. 2002
Denizli	Unreported	Erdem and Çanakçıoğlu 1977, Çanakçıoğlu 1983, Çanakçıoğlu and Mol 1998, Özdikmen 2008, Özdikmen et al. 2012
Denizli	Tavas	Gül-Zümreoğlu 1972
Düzce	Unreported	Özdikmen 2007, 2008; Özdikmen et al. 2012
Hatay	Unreported	Özdikmen 2008, Özdikmen et al. 2012
Hatay	Akbez	Pic 1892
Hatay	Central–Iskenderun (Cırtıman)	Tozlu et al. 2002
İçel	Unreported	Özdikmen 2008, Özdikmen et al. 2012

**Table 1. Continued.**

Province	Locality	Citations
İçel	Erdemli (Karahasanlı village)	Özdikmen 2006
İçel	Erdemli–Tarsus	Tozlu et al. 2002
İçel	Erdemli, Karahıdırlı	Özdikmen 2011, Cihan 2013, Cihan et al. 2013
İçel	Silifke	Özdikmen 2011, Cihan 2013, Cihan et al. 2013
İçel	Mut, Alahan	Özdikmen 2011, Cihan 2013, Cihan et al. 2013
İçel	Tarsus, Ayvalı	Özdikmen 2011, Cihan 2013, Cihan et al. 2013
İçel	Tarsus, Karabucak	Özdikmen 2011, Cihan 2013, Cihan et al. 2013
İçel	Tarsus, Yenice highway	Özdikmen 2011, Cihan 2013, Cihan et al. 2013
İçel	Çamlıyayala, İledin top	Özdikmen 2011, Cihan 2013, Cihan et al. 2013
İçel	Karakoyak	Özdikmen 2011, Cihan 2013, Cihan et al. 2013
İstanbul	Unreported	Erdem and Çanakçıoğlu 1977, Çanakçıoğlu 1983, Çanakçıoğlu and Mol 1998, Özdikmen 2008, Özdikmen et al. 2012
İstanbul	Polonez village	Acatay 1971, Tozlu et al. 2002
İzmir	Unreported	Erdem and Çanakçıoğlu 1977, Çanakçıoğlu 1983, Çanakçıoğlu and Mol 1998, Özdikmen 2008, Özdikmen et al. 2012
İzmir	Bornova	Gül-Zümreoğlu 1972
İzmir	Bornova–Kemalpaşa	Gül-Zümreoğlu 1975
Kahramanmaraş	Unreported	Kanat 1998, Özdikmen 2008, Özdikmen et al. 2012
Konya	Unreported	Özdikmen 2008, Özdikmen et al. 2012

**Table 1. Continued.**

Province	Locality	Citations
Konya	Akşehir	Tozlu et al. 2002
Muğla	Unreported	Erdem and Çanakçıoğlu 1977, Çanakçıoğlu 1983, Çanakçıoğlu and Mol 1998, Özdikmen 2008, Özdikmen et al. 2012, Lodos 1998
Muğla	Fethiye	Acatay 1971
Muğla	Marmaris	Öymen 1987
Osmaniye	Unreported	Özdikmen 2008, Özdikmen et al. 2012
Osmaniye	Çona village, Issizca village Karaçay	Agras 2006, Bahadıroğlu et al. 2009
Osmaniye	Çona village, Bahçe Road	Güven 2007, Özdikmen and Turgut 2009, Özdikmen et al. 2010
Osmaniye	Kadirli (Kabayar)	Tozlu et al. 2002
Turkey	Thrace, European	Althoff and Danilevsky 1997, Özdikmen 2008, Özdikmen et al. 2012
Turkey		Danilevsky and Miroshnikov 1985, Önder et al. 1987, Miroshnikov 1998

### Materials and Methods

A comprehensive review of the scientific literature was conducted to delineate the known distribution of *R. serricollis*. Host plants and ecological habits were recorded when available. Additional surveys for this insect were conducted by many researchers and are reported herein. We included information and data that are important in assessing the level of threat to the species. These protocols included geographic range, population data, and habitat preferences (International Union for Conservation of Nature 2012, Nieto and Alexander 2010).

### Results and Discussion

**Reported occurrence in Turkey.** *Rhaesus serricollis* is reported as occurring in 14 of Turkey's 81 provinces (Fig. 1). These are Adana, Antalya, Bilecik, Burdur,

**Table 2. Reported global occurrence of *Rhaesus serricollis*, with bibliographic citations.**

Reported Locale	Citations
Albania	Danilevsky 2007, Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Rapuzzi and Sama 2012, Kovács and Merkl 2013
Armenia	Özdikmen 2014
Azerbaijan	Özdikmen 2014
Balkan Peninsula	Heyrovský 1948, Sama 1993, Althoff and Danilevsky 1997, Sama et al. 2010
Bulgaria	Danilevsky 2007, Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Heyrovský 1948, Sama 1993, Althoff and Danilevsky 1997, Sama et al. 2010
Bulgaria (Asenovgrad)	Georgiev et al. 2006
Bulgaria (Lyubimets)	Georgiev and Doychev 2010
Caucasus Region	Demelt 1963, Villiers 1967, Agras 2006, Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Heyrovský 1948, Sama 1993, Althoff and Danilevsky 1997, Sama et al. 2010, Rapuzzi and Sama 2012, Cihan 2013, Özdikmen 2014
Cyprus	Heyrovský 1948, Sama 1993, Althoff and Danilevsky 1997, Sama et al. 2010, Özdikmen 2014
Dalmatia	Heyrovský 1948, Sama 1993, Althoff and Danilevsky 1997, Sama et al. 2010
East Mediterranean Region	Buse et al. 2013
Europe	Güven 2007, Özdikmen 2007, Marazzi 2009, Özdikmen and Turgut 2009
Europe (Balkans)	Cihan 2013
Europe (Southeast)	Özdikmen 2014
Georgia	Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Rapuzzi and Sama 2012, Özdikmen 2014
Germany	Demelt 1963, Villiers 1967, Agras 2006

**Table 2. Continued.**

Reported Locale	Citations
Greece	Danilevsky 2007, Demelt 1963, Villiers 1967, Atras 2006, Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Heyrovský 1948, Sama 1993, Althoff and Danilevsky 1997, Sama et al. 2010
Iran	Demelt 1963, Villiers 1967, Atras 2006, Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Heyrovský 1948, Sama 1993, Althoff and Danilevsky 1997, Sama et al. 2010, Rapuzzi and Sama 2012, Cihan 2013, Özdikmen 2014
Iran (East Azerbaijan province)	Sakenin et al. 2011
Iran (Northern, Western provinces)	Sama et al. 2008
Israel	Heyrovský 1948, Sama 1993, Althoff and Danilevsky 1997, Sama et al. 2010, Özdikmen 2014
Lebanon	Özdikmen 2014
Macedonia	Danilevsky 2007, Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Kovács and Merkl 2013
Near East Region	Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Cihan 2013
North Africa (Egypt)	Özdikmen 2014
Serbia	Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009
Syria	Demelt 1963, Villiers 1967, Atras 2006, Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Heyrovský 1948, Sama 1993, Althoff and Danilevsky 1997, Sama et al. 2010, Cihan 2013, Özdikmen 2014
Transcaucasus Region	Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Cihan 2013, Özdikmen 2014
Turkey	Danilevsky 2007, Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009, Heyrovský 1948, Sama 1993, Althoff and Danilevsky 1997, Sama et al. 2010, Cihan 2013, Özdikmen 2014



**Table 2. Continued.**

Reported Locale	Citations
Turkey (European)	Güven 2007, Özdikmen 2007, Özdikmen and Turgut 2009
Yugoslavia	Danilevsky 2007

Denizli, Düzce, Hatay, İçel, İstanbul, İzmir, Kahramanmaraş, Konya, Muğla, and Osmaniye provinces. It also has been recorded from the Toros Mountains. Location reports along with specific citation(s) of those reports are listed in Table 1.

**Reported occurrence outside Turkey.** *Rhaesus serricollis* is recorded from the Balkan peninsula from Dalmatia to southern Greece, Europe (Albania, Serbia, Macedonia, Greece, Bulgaria, European Turkey), Southeast Europe, Turkey, Caucasus, Transcaucasia (Armenia, Azerbaijan, Georgia), Iran, Syria, Cyprus, Israel, the Near East, Lebanon, and North Africa. It also has been introduced into Egypt (Özdikmen 2014). Citations of confirmed occurrence of *R. serricollis* are listed in Table 2, and the recorded distribution is shown in Fig. 1.

**Host plants.** *Rhaesus serricollis* is apparently polyphagous and develops in dead parts of living deciduous trees such as *Platanus orientalis* L. (Platanaceae), *Ficus* (Moraceae), *Quercus coccifera* subsp. *calliprinos* (Webb) Holmboe, *Q. ithaburensis* (Decne) Boissier, and *Populus alba* L. (Buse et al. 2013; Bytinski-Salz 1956; Bytinski-Salz and Sternlicht 1967; Georgiev and Doychev 2010; Halperin and Holzschuh 1993; Sama et al. 2008, 2010). Cihan (2013) and Özdikmen (2013) reported that the species lives on deciduous forest trees, including those of the genera *Fagus*, *Celtis*, *Platanus*, *Quercus*, *Castanea*, *Tilia*, *Juglans*, *Salix*, *Morus*, *Liquidambar*, and especially the conifers (*Pinus*). Specimens from Turkey were found on or in the deciduous species *Juglans regia* L., *Liquidambar orientalis* L., *Platanus orientalis*, *Morus alba* L., and the conifers *Pinus brutia* Steven and *P. nigra* Arnold either as adults or larvae (Acatay 1971, Adlbauer 1988, Bahadıroğlu et al. 2009, Bense 1995, Çanakçıoğlu 1983, Demelt 1963, Erdem and Çanakçıoğlu 1977, Gül-Zümreoğlu 1975, Hoskovec and Rejzek 2012, Öymen 1987, Svacha and Danilevsky 1987). *Rhaesus serricollis* also is the only long-horned beetle species reported from Oriental sweet gum, *L. orientalis*, in Turkey (Acatay 1971, Çanakçıoğlu 1983, Cebeci and Özdikmen 2010, Erdem and Çanakçıoğlu 1977, Öymen 1987).

**Life cycle and biology.** *Rhaesus serricollis* larvae live in rotting and wet wood (Agras 2006, Gül-Zümreoğlu 1975). Adults and larvae can be collected only from the host plants growing in lowland habitats between 5 and 1,000 m above sea level. Duration of the life cycle is at least 3 yr. The overwintering stage is most likely the larval stage. Larvae live in rotten wood of large trunks (both standing or fallen) of the host plants. Pupation occurs in the wood in the spring and summer. Adults are crepuscular, nocturnal, and are attracted to light. They actively fly in early spring and summer (between May and August) (Özdikmen 2013).

**Status and conservation of threatened species.** Nieto and Alexander (2010) report that 13% of the species of saproxylic beetles are considered Near

Threatened in Europe. *Rhaesus serricollis* is included in those 56 species. According to Özdikmen (2014), *R. serricollis* is probably widely distributed in Turkey and suggested that the species be listed in the category of Least Concern in the Turkish Red List. Based on its distribution, collection dates, and records from Turkey, we concur with that placement.

### References Cited

- Acatay, A. 1971.** Über das Auftreten einiger Forstschädlingen in der Türkei. Anz. Schaedlingskd., Pflanz.-Umweltschutz 11: 162–165.
- Aldbauer, K. 1988.** Neues zur Taxonomie und Faunistik der Bockkäferfauna der Türkei (Coleoptera, Cerambycidae). Entomofauna 9: 257–297.
- Agras, M. 2006.** Amanos Dağı (Osmaniye İli) Cerambycidae ve Buprestidae (Coleoptera) familyalarına ait bazı böcek türleri ve yükseltiyeye göre dağılımı üzerine Araştırmalar. Yüksek Lisans Tezi. Kahramanmaraş Sütçü İmam Üniversitesi Biyoloji Anabilim Dalı. 41 pp.
- Althoff, J. and M.L. Danilevsky. 1997.** A check-list of longicorn beetles (Coleoptera, Cerambycoidea) of Europe. Slovensko entomolosko drustvo Stefana Michielija, Ljubljana. 64 pp.
- Avcı, M., O. Sarıkaya, N. Jansson and M. Coşkun. 2010.** The beetle fauna on old oaks (*Quercus* spp.) in Kasnak Forest east of Isparta in Turkey, Pp. 126. In The Oak—Ecology, History, Management and Planning II. 2010 June 1–3; Suleyman Demirel Univ., Isparta, Turkey.
- Bahadıroğlu, C., M. Agras and Ü.F. Salman. 2009.** Amanos Dağı (Osmaniye İli) Cerambycidae Familyasına ait türler ve yükseltiyeye göre dağılımları üzerine araştırmalar. KSÜ Doğa Bil. Derg. 12(1): 1–8.
- Bense, U. 1995.** Longhorn Beetles Illustrated Key to the Cerambycidae and Vesperidae of Europe. Druckerei Steinmeier, Nördlingen, Germany. 512 pp.
- Bodemeyer, H.E.V. 1906.** Beiträge zur Käferfauna von Klein Asien. Dtsch. Entomol. Z. 2: 417–437.
- Buse, J., T. Assmann, A.L.L. Friedman, O. Rittner and T. Pavlicek. 2013.** Wood-inhabiting beetles (Coleoptera) associated with oaks in a global biodiversity hotspot: A case study and checklist for Israel. Insect Conserv. Diversity 6: 687–703.
- Bytinski-Salz, H. 1956.** The Cerambycidae of Israel. Bull. Res. Council. Isr. 5B: 207–226.
- Bytinski-Salz, H. and M. Sternlicht. 1967.** Insects associated with oaks (*Quercus*) in Israel. Isr. J. Entomol. 2: 107–143.
- Çanakçıoğlu, H. 1983.** Orman entomolojisi: Özel bölüm. İstanbul Univ., Orman Fak. Yayını, İstanbul, 349: 535.
- Çanakçıoğlu, H. and T. Mol. 1998.** Orman entomolojisi: Zararlı ve yararlı böcekler. İstanbul Univ., Orman Fak. Yayınları 451: 144–174.
- Cebeci, H. and H. Özdikmen. 2010.** A synopsis on the genus *Rhamnusium* Latreille, 1829 with a new record (Coleoptera: Cerambycidae) for Anatolian fauna from a new host plant, *Liquidambar orientalis* Miller (Hamamelidaceae). Munis Entomol. Zool. 5: 131–139.
- Cihan, N. 2013.** Doğu Akdeniz Ormancılık Araştırma Enstitüsü Entomoloji Müzesindeki Teke Böcekleri (Coleoptera: Cerambycidae) üzerine PRIONINAE, LEPTURINAE, DORCADIONINAE VE LAMIINAE itibarıyla taksonomik ve faunistik. Yüksek Lisans Tezi. Gazi Üniversitesi Fen Bilimleri Enstitüsü, Ankara, Turkey. 184 pp.
- Cihan, N., H. Özdikmen and F. Aytar. 2013.** Contributions of the longhorned beetles knowledge of Turkey by the subfamilies Prioninae, Lepturinae, Dorcadioninae and Lamiinae (Coleoptera: Cerambycidae). Munis Entomol. Zool. 8: 883–894.
- Coşkun, M., N. Jansson, M. Avcı and O. Sarıkaya. 2010.** Rich and unique beetle fauna found on pollarded (coppice) oaks (*Quercus* spp.) in southern Turkey, Pp. 120. In The

- Oak—Ecology, History, Management and Planning II. 2010 June 1–3; Suleyman Demirel Univ., Isparta, Turkey.
- Danilevsky, M.L. 2007.** A check-list of longicorn beetles (Coleoptera, Cerambycoidea) of Europe. A.N. Severtsov Institute of Ecology and Evolution, Moscow. Revised 1 November 2007, accessed 29 January 2013. (<http://www.cerambycidae.net>).
- Danilevsky, M.L. and A.I. Miroshnikov. 1985.** Timber-beetles of Caucasus (Coleoptera, Cerambycidae), A Key Book. Krasnodar Publ., Krasnodar Territory, Russia. 419 pp.
- Demelt, C.V. 1963.** Beitrag zur Kenntnis der Cerambycidenfauna Kleinasiens und 13. Beitrag zur Biologie palaearkt. Cerambyciden, sowie Beschreibung einer neuen Oberea-Art. Entomol. Bl. 59: 132–151.
- Erdem, R. and H. Çanakçıoğlu. 1977.** Türkiye odun zararlıları, Pp. 113–134. İstanbul Üniversitesi Orman Fakültesi Yayını, İstanbul.
- Georgiev, G.T. and D. Doychev. 2010.** New xylophagous beetles (Insecta: Coleoptera) on poplars in Bulgaria. Acta Zool. Bulg. 62: 175–180.
- Georgiev, G., E. Migliaccio and D. Doychev. 2006.** Longhorn beetles (Coleoptera: Cerambycidae) in Western Rhodopes (Bulgaria), Pp. 347–360. In Beron, P. (ed.), Biodiversity of Bulgaria. 2. Biodiversity of Western Rhodopes (Bulgaria and Greece). Pensoft Publishers, Sofia, Bulgaria.
- Gücü, A.C. and F. Erkan. 2005.** A holistic approach for the conservation of the Mediterranean monk seal on the Cilician coast of Turkey. In INOC International Workshop on Marine and Coastal Protected Areas; 2005 Mar 23–25; Meknes, Morocco. 11 pp.
- Gül-Zümreöğlu, S. 1972.** Böcek ve Genel Zararlılar Kataloğu (1928–1969). T.C. Tarım Bakanlığı Yayınları, Bornova, İzmir. 119 pp.
- Gül-Zümreöğlu, S. 1975.** Ege Bölgesi Cerambycidae Türleri, Taksonomileri Konukçuları ve Yayılış Alanları Üzerinde Araştırmalar. T.C. Gıda-Tarım ve Hayvancılık Bakanlığı Zirai Mücadele ve Zirai Karantina Genel Müdürlüğü Araştırma Eserleri Serisi Yay. 28. İstiklal Matbaası, İzmir. 208 pp.
- Gürkan, T., E. Atay and N. Jansson. 2010.** The beetle fauna on old oaks (*Quercus* spp) in a cemetery in Adana Region, South Turkey, Pp. 122. In The Oak—Ecology, History, Management and Planning II. 2010 June 1–3; Suleyman Demirel Univ., Isparta, Turkey.
- Güven, M. 2007.** Batı Toroslar ve Güneydoğu Toroslar PRIONINAE-CERAMBYCINAE (Coleoptera: Cerambycidae) faunaları üzerine taksonomik, sistematik ve zoocoğrafik araştırmalar. Yüksek Lisans Tezi. Gazi Üniversitesi Fen Bilimleri Enstitüsü, Ankara, Turkey. 192 pp.
- Halperin, J. and C. Holzschuh. 1993.** Host plant of Israeli Cerambycidae with new records. Phytoparasitica 21: 23–37.
- Heyrovský, L. 1948.** Contribution à la faune des Longicornes de la Palestine. Cas. Cesk. Spol. Entomol. 45(1–2): 17–22.
- Hoskovec, M. and M. Rejzek. 2012.** Longhorn beetles (Cerambycidae) of the West Palearctic Region. 31 October 2014. (<http://www.cerambyx.uochb.cz>).
- International Union for Conservation of Nature. 2012.** International Union for Conservation of Nature Red List of Threatened Species. Version 2012. 2.0. 30 March 2013. ([www.iucnredlist.org](http://www.iucnredlist.org)).
- Kahraman, A., M. Önder and E. Ceyhan. 2011.** Biodiversity and biosecurity in Turkey. International Conference on Biology, Environment and Chemistry, IPCBEE. IACSIT Press, Singapore. Int. Proc. Chem., Biol. Environ. Eng. 24: 33–37.
- Kanat, M. 1998.** Kahramanmaraş ormanlarında önemli zararlı böceklerin araştırılması. PhD Dissertation. Univ. Karadeniz Teknik, Trabzon, Turkey. 127 pp.
- Koçak, A.Ö. and M. Kemal. 2009.** List of the coleopteran genera and species recorded in Turkey based upon the info-system of Cesa. Cent. Entomol. Stud. Ankara 53: 1–213.
- Kovács, T. and O. Merkl. 2013.** Beetles from Albania, Macedonia and Montenegro, with new country records (Coleoptera). Folia Hist. Nat. Mus. Matraensis 37: 89–92.

- Lodos, N. 1998.** Türkiye Entomolojisi VI (Genel, Uygulamalı ve Faunistik). Ege Üniversitesi Ziraat Facultesi Yayınları 529. 300 pp.
- Marazzi, G. 2009.** Checklist of the Prioninae. Checklist of the Cerambycidae (Subfamily Prioninae) of the world. 29 May 2014. (<http://marazziprioninae.com/wp-content/uploads/2014/02/CHECKLIST-2009-Marazzi-Collection.pdf>).
- Miroschnikov, A.I. 1998.** *Rhaesus* Motschulsky, 1875—The valid name of the genus incorrectly called *Rhesus* Motschulsky, 1838 (Coleoptera, Cerambycidae). Rev. Entomol. 77: 616–617.
- Nieto, A. and K.N.A. Alexander. 2010.** European Red List of Saproxyllic Beetles. Publications Office of the European Union, Luxembourg. 39 pp.
- Önder, F., Y. Karsavuran, S. Tezcan and P. Önder. 1987.** Türkiye’de tarım, orman ve evcil hayvanlarda hayvansal kökenli zararlı ve yaralı türlerin bilimsel ve Türkçe isimleri. T.C. Tar. Orm. Köyişl. Bakanlığı, Mesleki Yayınlar 8: 1–120.
- Öymen, T. 1987.** The Forest Cerambycidae of Turkey. Istanbul University of Forestry, İstanbul. 146 pp.
- Özdkimen, H. 2006.** Contribution to the knowledge of Turkish longicorn beetles fauna (Coleoptera: Cerambycidae). Munis Entomol. Zool. 1: 71–90.
- Özdkimen, H. 2007.** The longhorn beetles of Turkey (Coleoptera: Cerambycidae). Part I—Black Sea Region. Munis Entomol. Zool. 2: 179–422.
- Özdkimen, H. 2008.** The longicorn beetles of Turkey (Coleoptera: Cerambycidae). Part III—Aegean Region. Munis Entomol. Zool. 3: 355–436.
- Özdkimen, H. 2011.** The longicorn beetles of Turkey (Coleoptera: Cerambycidae). Part IV—Mediterranean Region. Munis Entomol. Zool. 6: 6–145.
- Özdkimen, H. 2013.** Review on the biology of Turkish Cerambycoidea (Coleoptera). Part I—Vesperidae and Cerambycidae (Prioninae). Munis Entomol. Zool. 8: 278–284.
- Özdkimen, H. 2014.** Turkish Red List categories of longicorn beetles (Coleoptera: Cerambycidae). Part I—Subfamilies Vesperinae and Prioninae. Munis Entomol. Zool. 9: 245–253.
- Özdkimen, H., M. Güven and C. Gören. 2010.** Longhorned beetles fauna of Amanos Mountains, Southern Turkey (Coleoptera: Cerambycidae). Munis Entomol. Zool. 5: 1141–1167.
- Özdkimen, H., N. Mercan and H. Tunç. 2012.** Longhorned beetles of Düzce province in Turkey (Coleoptera: Cerambycidae). Munis Entomol. Zool. 7: 714–731.
- Özdkimen, H. and S. Turgut. 2009.** A synopsis of Turkish Vesperinae Mulsant, 1839 and Prioninae Latreille, 1802 (Coleoptera: Cerambycidae). Munis Entomol. Zool. 4: 402–423.
- Pic, M. 1892.** Séance du 14 Décembre 1892 [note]. Ann. Soc. Entomol. Fr. 61: CCUXCCLX.
- Rapuzzi, P. and G. Sama. 2012.** Contributo alla conoscenza dei Cerambycidae di Albania (Coleoptera, Cerambycidae). Atti Mus. Civ. Stor. Nat. Trieste 55: 181–234.
- Sakenin, H., N. Samin, S. Moemen Beitollahi, S. Ezzatpanah, M. Havaskary, J. Rastegar, A. Valizadeh and M.J. Shakouri. 2011.** A study on the longhorn beetles (Coleoptera: Cerambycidae) from north-western Iran. Calodema 143: 1–19.
- Sama, G. 1993.** Notes on *Neomusaria* Plavilstshikov, 1928 with description of a new species from Turkey. Koleopterol. Rundsch. 63: 293–297.
- Sama, G., J. Buse, E. Orbach, A.L.L. Friedman, O. Rittner and V. Chikatunov. 2010.** A new catalogue of the Cerambycidae (Coleoptera) of Israel with notes on their distribution and host plants. Munis Entomol. Zool. 5: 1–51.
- Sama, G., N. Seddighi and A.A. Talebi. 2008.** Preliminary note for a checklist of the Cerambycidae of Iran (Coleoptera-Cerambycidae). Biocosme Méditerranéenne Nice 25: 101–126.
- Şekercioğlu, Ç.H., S. Anderson, E. Akcay, R. Bilgin, Ö.E. Can, G. Semiz, Ç. Tavşanoğlu, M.B. Yokeş, A. Soyumert, K. İpekdal, İ.K. Sağlam, M. Yücel and H.N. Dalfes. 2011.** Turkey’s globally important biodiversity in crisis. Biol. Conserv. 144: 2752–2769.
- Sundseth, K. and E.L.T.D. Brussels. 2009.** Natura 2000 in the Mediterranean Region. Office for Official Publications of the European Communities, Luxembourg. 12 pp.

- Svacha, P. and M.L. Danilevsky. 1987.** Cerambycoid larvae of Europe and Soviet Union (Coleoptera: Cerambycoidea). Part I. Acta Univ. Carol., Biol. 30: 1–186.
- Tozlu, G., M. Rejzek and H. Özbek. 2002.** A contribution to the knowledge of Cerambycidae (Coleoptera) fauna of Turkey. Part I: Subfamilies Prioninae to Cerambycinae. Biocosme Mésogéen Nice 19(1–2): 55–94.
- Villiers, A. 1967.** Coléoptères Cérambycides de Turquie (1. Partie). L'Entomologiste 23(1): 18–22.
- Vlachogianni, T., M. Vogrin and M. Scoullou. 2012.** Biodiversity in the Mediterranean region. MIO-ECSDE 2012. Mediterranean Information Office for Environment, Culture and Sustainable Development. Federation of Mediterranean Non-Governmental Organisations for Environment and Sustainable Development. Athens, Greece. 19 pp.