

505. PHYTOECIA *Faldermanni* DEJEAN. Tab. X. f. 3.

Nigra, dense pubescens, opaca; vertice vittis duabus latis rufis; antennis, thorace, pedibusque obscure rufescentibus, nigro-maculatis; thorace tuberculis septem nigris; elytris valde depressis, lurido-flavis; lateribus nigris, glabris; ano rufo-testaceo.

Longit. 5 lin. Lat. $4\frac{2}{5}$ lin.

Statura et colore Saperdæ Argo Panz. valde affinis et similis, tamen dimidio fere minor præsertim angustior.

Caput rotundatum, convexum, nigrum, opacum, punctis grossis sub-confluentibus dense adpersum, basi parce antice densius testaceo-pubescens sub-villosum, vertice vittis duabus latis rufis signatum ibique tenuiter canaliculatum; ore pilis pallidis rigidis dense obsito. Oculi parum prominuli, aterrimi, postice rufo-cingulati. Antennæ corpore multo breviores, pube tenuissima grisea et pilis rigidis nigris vage immixtis obductæ; articulo primo nigro, creberrime punctulato; 2-4 obscure rufescentibus apice nigris; subsequentibus obscure fuscis. Thorax longitudine paullo latior, cylindricus, basi apiceque truncatus, lateribus in medio vix dilatatus, undique crebre punctatus, dense testaceo-villosus, obscure rufescens, opacus, intra apicem transversim late impressus, in disco tuberculis septem 4, 3 nigris nitidis notatus. Scutellum transversum, apice obtuse rotundatum, supra planum, densissime villosum. Elytra thorace dimidio fere latiora, basi truncata, apicem versus oblique valde attenuata, apice sub-dehiscentia, truncata, supra valde depressa, plana, minute vix visibiliter remote punctata, pube lurido-flava densissima et valde depressa ves-

tita, lateribus declivia ibique nuda, nigra, rude et crebre punctata, carinato-marginata, margine sub humero rufo-testaceo; præterea in medio disci singulo elytri canalicula longitudinali valde obsoleta instructa. Corpus subtus nigrum, opacum, densissime cano-pubescentis, segmentis duobus ulterioribus rufo-testaceis, postice in medio late nigro-marginatis. Pedes breviusculi, dense pubescentes, obscure rufescentes, femorum basi nonnihil infuscati sed femorum tiliarumque apicibus nigris; tarsi nigris subtus densissime fusco-spongiosis.

506. PHYTOECIA *diademata* MIHL. Tab. X. f. 3.

Brevis, parallela, atra, sub-glabra; capite, thorace pedibusque luteis, nigro-maculatis; antennis bi-coloribus; elytris testaceis, disco obsolete bicarinatis, apice late nigris.

Longit. $4\frac{1}{5}$ lin. Lat. $1\frac{2}{5}$ lin.

Statura et fere Longitudo Saperdæ prætextatæ Stev. tamen triplo fere latior.

Caput magnum, crassum, sub-globosum, luteum, parce pilosum, sub-nitidum, ubique confertissime at obsolete punctulatum, lineola angustissima longitudinali, vertice in medio vix elevata et antice impressa instructum, ante oculos superne macula utrinque et vitta frontali nigris signatum, intra apicem linea angusta transversim impressum; fronte convexa. Oculi modice prominuli, aterrimi. Antennæ corporis dimidio paullo longiores, lineares, crassiusculæ, nigræ, latere interiori testaceæ; articulo basali luteo, superne vitta angusta nigra notato. Thorax brevissimus, longitudine plus duplo latior, supra luteus, subtus niger, basi apiceque truncatus, lateribus valde



Phytoecia puncticollis Menétr. 2. *P. azurea* Steven. 3. *P. Faldermanni*
 Dej. 4. *P. diademata* Fald. 5. *P. piceosa* Fald. 6. *Agapanthia smaragdina*
 Dej. 7. *A. chalybea* Fald. 8. *Phagium fasciculatum* Fald. 9. *Toxotus in-*
sitivus Boeber. 10, 11. *Toxotus persicus* ♂, ♀. Fald.